

The line bundles on the moduli of parabolic G-bundles over curves and their sections

Yves Laszlo and Christoph Sorger

Abstract

Let X be a smooth, complete and connected curve and G be a simple and simply connected algebraic group over $\mathbb{C}$. We calculate the Picard group of the moduli stack of quasi-parabolic G -bundles and identify the spaces of sections of its members to the conformal blocs of Tsuchiya, Ueno and Yamada. We describe the canonical sheaf on these stacks and show that they admit a unique square root, which we will construct explicitly. Finally we show how the results on the stacks apply to the coarse moduli spaces and recover (and extend) the Drezet-Narasimhan theorem. We show moreover that the coarse moduli spaces of semi-stable SO_r -bundles are not locally factorial for $r \geq 7$.

1. Introduction.

(1.1) Fix a simple and simply connected algebraic group G over $k = \mathbb{C}$ and a Borel subgroup $B \subset G$. Let X be a smooth, complete and connected curve over k and $p_1, \dots, p_n$ be distinct points of X , labeled by standard (*i.e.* containing B) parabolic subgroups $P_1, \dots, P_n$ of G (we allow $n = 0$). Let $\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P})$ be the moduli stack of quasi-parabolic G -bundles of type $\underline{P} = (P_1, \dots, P_n)$ at $\underline{p} = (p_1, \dots, p_n)$ and denote by $X(P_i)$ the character group of P_i .

THEOREM.— *There is $\mathcal{L} \in \text{Pic}(\text{Bun}_G)$ such that we have an isomorphism*

$$\gamma : \text{Pic}(\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P})) \xrightarrow{\sim} \mathbb{Z}\mathcal{L} \times \prod_{i=1}^n X(P_i).$$

If G is of type A or C, then $\mathcal{L}$ is the determinant of cohomology (cf. 7.1). If G is of type B, D or G_2 , then $\mathcal{L}$ is the pfaffian of cohomology (cf. 7.2).

If G is of type E_6, E_7, E_8, F_4 we believe that we have $\mathcal{L}^{\otimes d(G)} = \mathcal{D}_{\rho(G)}$ where respectively $d(G) = 6, 12, 60, 6$ and $\mathcal{D}_{\rho(G)}$ is the determinant of cohomology (cf. 7.1) associated to the fundamental representation $\rho(G) = \varpi_6, \varpi_7, \varpi_8, \varpi_4$ (cf. the discussion in 1.4).

(1.2) Suppose that the points $p_1, \dots, p_n$ are instead labeled by finite dimensional simple representations $\lambda_1, \dots, \lambda_n$ of G and that an additional integer ℓ , the *level*, is fixed. The choice of a representation λ of G is equivalent to the choice of a standard parabolic subgroup $P \subset G$ and a character $\chi \in X(P)$. Therefore, the labeling of the points $p_1, \dots, p_n$ by the representations $\lambda_1, \dots, \lambda_n$ defines the type $\underline{P}$ of a quasi-parabolic G -bundle, that is the stack $\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P})$ and, by the above theorem, a line bundle $\mathcal{L}(\ell, \underline{\chi})$ over $\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P})$. The global sections of $\mathcal{L}(\ell, \underline{\chi})$ give a vector space, the space of *generalized parabolic G -theta-functions of level ℓ* , which is canonically associated to $(X, \underline{p}, \underline{\lambda})$. In mathematical physics, the rational conformal field theory of Tsuchiya, Ueno and Yamada [32] associates also to $(X, \underline{p}, \underline{\lambda}, \ell)$ a vector space: the space of *conformal blocks* $V_X(\underline{p}, \underline{\lambda}, \ell)$ (cf. [31] for an overview).

THEOREM.— Suppose that G is classical or G_2 . Then there is a canonical isomorphism

$$H^0(\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P}), \mathcal{L}(\ell, \underline{\chi})) \xrightarrow{\sim} V_X(\underline{p}, \underline{\lambda}, \ell). \quad (1.2.1)$$

In particular, $\dim H^0(\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P}), \mathcal{L}(\ell, \underline{\chi}))$ is given by the Verlinde dimension formula.

For $n = 0$, this has been proved independently by Beauville and the first author [3] for $G = \text{SL}_r$ and by Faltings [10] and Kumar, Narasimhan and Ramanathan [17] for arbitrary simple and simply connected G . For arbitrary n and $G = \text{SL}_r$ this has been proved by Pauly [24] and can be proved for arbitrary simple and simply connected G using (1.3) (and therefore [8]) and (5.1) below following the lines of [3] and [24]. This is the subject of Section 10.

(1.3) The above results are proved via the *uniformization* theorem: restrict for simplicity of the introduction to $n = 0$. Suppose $p \in X$ and denote $X^* = X - p$. Define $D = \text{Spec}(\hat{\mathcal{O}}_p)$, where $\hat{\mathcal{O}}_p$ is the formal completion of the local ring $\mathcal{O}_p$ at p and $D^* = \text{Spec}(K_p)$ where K_p is the quotient field of $\hat{\mathcal{O}}_p$. Let L_G (resp. L_G^+ , resp. L_G^X) be the group of algebraic morphisms from D^* (resp. D , resp. X^*) to G .

THEOREM.— The algebraic stack Bun_G is canonically isomorphic to the double quotient stack $L_G^X \backslash L_G / L_G^+$. Moreover, the projection map

$$\pi : \mathcal{Q}_G := L_G / L_G^+ \rightarrow \text{Bun}_G$$

is locally trivial for the étale topology.

This is proved in [3] for $G = \text{SL}_r$. The extension to arbitrary G has been made possible by Drinfeld and Simpson [8] in response to a question by the first author. They prove that if S is a k -scheme and E a G -bundle over $X \times S$ then, locally for the étale topology on S , the

restriction of E to $X^* \times S$ is trivial, which is essential for the proof. The above theorem is valid more generally for semi-simple G . In characteristic p , one has to replace “étale” by “fppf” if p divides the order of $\pi_1(G(\mathbb{C}))$.

(1.4) Consider the pullback morphism

$$\pi^* : \text{Pic}(\text{Bun}_G) \longrightarrow \text{Pic}(\mathcal{Q}_G).$$

The Picard group of $\mathcal{Q}_G$ is known ([21],[17]) to be canonically isomorphic to $\mathbb{Z}$, which reduces proving Theorem 1.1 to proving that π^* is an isomorphism. We will show that the injectivity of α will follow from the fact that L_G^X has no characters which in turn will follow from the fact that L_G^X is reduced and connected. Moreover, the surjectivity of α would follow from the simple connectedness of L_G^X . Both topological properties, connectedness and simple connectedness of L_G^X are affirmed in [17] and we believe them to be true. Whereas we will prove the connectedness of L_G^X , following an idea of V. Drinfeld, we do not see how to prove the simple connectedness of L_G^X . The injectivity is enough to prove the first part of Theorem 1.1, but to identify the generator $\mathcal{L}$ we should prove the surjectivity of α . For classical G and G_2 we do this by constructing a line bundle on Bun_G pulling back to the generator of $\text{Pic}(\mathcal{Q}_G)$.

(1.5) The pfaffian construction (7.2) may be used to prove the following, valid over k algebraically closed of characteristic $\neq 2$.

PROPOSITION.— *Suppose G is semi-simple. Then, for every theta-characteristic κ on X , there is a canonical square-root $\mathcal{P}_\kappa$ of the dualizing sheaf ω_{Bun_G} of Bun_G .*

(1.6) The last section will be devoted to Ramanathans moduli spaces M_G of semi-stable G -bundles. We will show how some of the results for the stack Bun_G will be true also for the moduli spaces M_G . In particular we will recover (and extend) the Drezet-Narasimhan theorem.

THEOREM.— *There is a canonical isomorphism $\text{Pic}(M_G) \xrightarrow{\sim} \mathbb{Z}L$. If G is of type A or C then L is the determinant bundle and moreover, in this case M_G is locally factorial. If G is of type B_r , D_r , $r \geq 4$ or G_2 then L or $L^{\otimes 2}$ is the determinant bundle.*

This theorem has also been proved, independently and with a different method, by Kumar and Narasimhan [18].

The question whether M_G is locally factorial for G of type other than the simply connected groups of type A, C is the subject of a forthcoming paper. We show there for example that $\text{Pic}(M_{\text{Spin}_r})$ is generated by the determinant of cohomology and in particular that M_{Spin_r} is not

locally factorial for $r \geq 7$ by “lifting” to Spin_r the proof we give here (11.7) for the analogous statement for M_{SO_r} .

We would like to thank A. Beauville and C. Simpson for useful discussions and V. Drinfeld for his suggestion in (5.1) and for pointing out an inaccuracy in an earlier version of this paper.

2. Some Lie theory.

Throughout this section k will be an algebraically closed field of characteristic zero.

(2.1) The general set up. Let $\mathfrak{g}$ be a simple finite dimensional Lie algebra over k . We fix a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ and denote by Δ the associated root system. We have the root decomposition $\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_\alpha$. The Lie subalgebra $\mathfrak{g}_{-\alpha} \oplus [\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] \oplus \mathfrak{g}_\alpha$, isomorphic as a Lie algebra to $\mathfrak{sl}_2$, will be denoted by $\mathfrak{sl}_2(\alpha)$. Moreover we choose a basis $\Pi = \{\alpha_1, \dots, \alpha_r\}$ of Δ and we denote by Δ_+ the set of positive roots (with respect to Π). Put $\mathfrak{b} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta_+} \mathfrak{g}_\alpha$. For each $\alpha \in \Delta_+$, we denote by H_α the coroot of α , *i.e.* the unique element of $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}]$ such that $\alpha(H_\alpha) = 2$, and we denote by $X_\alpha \in \mathfrak{g}_\alpha$ and $X_{-\alpha} \in \mathfrak{g}_{-\alpha}$ elements such that

$$[H_\alpha, X_\alpha] = 2X_\alpha, \quad [H_\alpha, X_{-\alpha}] = -2X_{-\alpha}, \quad [X_\alpha, X_{-\alpha}] = H_\alpha.$$

When α is one of the simple roots α_i , we write H_i, X_i, Y_i instead of $H_{\alpha_i}, X_{\alpha_i}, Y_{\alpha_i}$. Let (ϖ_i) be the basis of $\mathfrak{h}^*$ dual to the basis (H_i) . Denote by P the weight lattice and by $P_+ \subset P$ the set of dominant weights. Given a dominant weight λ , denote L_λ the associated simple $\mathfrak{g}$ -module with highest weight λ and v_λ its highest weight vector. Finally $(,)$ will be the Cartan-Killing form normalized such that for the highest root θ we have $(\theta, \theta) = 2$.

(2.2) Loop algebras. Let $L\mathfrak{g} = \mathfrak{g} \otimes_k k((z))$ be the *loop algebra* of $\mathfrak{g}$ and $L\mathfrak{g}^+ = \mathfrak{g} \otimes_k k[[z]]$ its subalgebra of *positive loops*. There is a natural 2-cocycle

$$\begin{aligned} \psi_{\mathfrak{g}} : \quad L\mathfrak{g} \quad \times \quad L\mathfrak{g} \quad &\longrightarrow \quad k \\ (X \otimes f, \quad Y \otimes g) \quad &\mapsto \quad (X, Y) \text{Res}(gdf) \end{aligned}$$

defining a central extension $\widehat{L\mathfrak{g}}$ of $\mathfrak{g}$:

$$0 \longrightarrow k \longrightarrow \widehat{L\mathfrak{g}} \longrightarrow L\mathfrak{g} \longrightarrow 0.$$

Every other cocycle is a scalar multiple of $\psi_{\mathfrak{g}}$ and the above central extension is universal. Let $\widehat{L\mathfrak{g}}^+$ be the extension of $L\mathfrak{g}^+$ obtained by restricting the above extension to $L\mathfrak{g}^+$. As the cocycle is trivial over $L\mathfrak{g}^+$ this extension splits.

Let ℓ be a positive integer. A representation of $\widehat{L}_{\mathfrak{g}}$ is of *level* ℓ if the center c acts by multiplication by ℓ . Such a representation is called integrable if $X \otimes f$ acts locally nilpotent for all $X \otimes f \in \mathfrak{g}_\alpha \otimes_k k((z))$. The theory of affine Lie algebras [13] affirms that the irreducible integrable representations of level ℓ of $\widehat{L}_{\mathfrak{g}}$ are classified (up to isomorphism) by the weights $P_\ell = \{\lambda \in P_+ / (\lambda, \theta) \leq \ell\}$. We denote by $\mathcal{H}_{\lambda, \ell}$ the irreducible integrable representation of level ℓ and highest weight $\lambda \in P_\ell$. If $\lambda = 0$, the corresponding representation, which we denote simply by $\mathcal{H}_\ell$, is called the *basic representation of level* ℓ .

(2.3) The Dynkin index. Let $\rho : \mathfrak{g} \rightarrow \mathfrak{sl}(V)$ be a representation of $\mathfrak{g}$. Then ρ induces a morphism of Lie algebras $L_{\mathfrak{g}} \rightarrow L_{\mathfrak{sl}(V)}$. Pull back the universal central extension to $L_{\mathfrak{g}}$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & k & \longrightarrow & \widetilde{L}_{\mathfrak{g}} & \longrightarrow & L_{\mathfrak{g}} \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & k & \longrightarrow & \widehat{L_{\mathfrak{sl}(V)}} & \longrightarrow & L_{\mathfrak{sl}(V)} \longrightarrow 0 \end{array}$$

The cocycle of the central extension $\widetilde{L}_{\mathfrak{g}}$ is of the type $d_\rho \psi_{\mathfrak{g}}$. Define the *Dynkin index* of the representation ρ of $\mathfrak{g}$ by the number d_ρ .

(2.4) LEMMA.— Let $V = \sum_\lambda n_\lambda e^\lambda$ be the formal character of V . Then we have

$$d_\rho = \frac{1}{2} \sum_\lambda n_\lambda \lambda(H_\theta)^2$$

Proof. By definition of the cocycle, we have $d_\rho = \text{Tr}(\rho(X_\theta)\rho(X_{-\theta}))$. Decompose the $\mathfrak{sl}_2(\theta)$ -module, V as $\oplus V^{(d_i)}$, where $V^{(d_i)}$ is the standard irreducible $\mathfrak{sl}_2$ -module with highest weight d_i . We may realize $V^{(d_i)}$ as the vector space of homogeneous polynomials in 2 variables x and y of degree d_i . Then X_θ acts as $x\partial/\partial y$, and $X_{-\theta}$ as $y\partial/\partial x$. Using the basis $x^l y^{d_i-l}$, $l = 0, \dots, d_i$ of $V^{(d_i)}$, we see

$$d_\rho = \sum_i \sum_{k=0}^{d_i} k(d_i + 1 - k).$$

The formal character of the $\mathfrak{sl}_2(\theta)$ -module $V^{(d)}$ is $\sum_{k=0}^d e^{d\rho_\theta - k\alpha_\theta}$ where α_θ is the positive root of $\mathfrak{sl}_2(\theta)$ and $\rho_\theta = \frac{1}{2}\alpha_\theta$. Therefore we are reduced to prove the equality

$$\sum_{k=0}^d k(d+1-k) = \frac{1}{2} \sum_{k=0}^d [d\rho_\theta - k\alpha_\theta](H_\theta)]^2 = \frac{1}{2} \sum_{k=0}^d [d-2k]^2$$

which is easy. $\square$

(2.5) REMARK.— The Dynkin index of a representation has been introduced to the theory of G -bundles over a curve by Faltings [10] and Kumar, Narasimhan, Ramanathan [17].

We are interested here in the *minimal Dynkin index* $d_{\mathfrak{g}}$ defined to be $\min d_{\rho}$ where ρ runs over all representations $\rho : \mathfrak{g} \rightarrow \mathfrak{sl}(V)$.

(2.6) PROPOSITION.— *The minimal Dynkin index $d_{\mathfrak{g}}$ is as follows*

Type of $\mathfrak{g}$	A_r	$B_r, r \geq 3$	C_r	$D_r, \geq 4$	E_6	E_7	E_8	F_4	G_2
$d_{\mathfrak{g}}$	1	2	1	2	6	12	60	6	2
λ s.t. $d_{\mathfrak{g}} = d_{\rho(\lambda)}$	ϖ_1	ϖ_1	ϖ_1	ϖ_1	ϖ_6	ϖ_7	ϖ_8	ϖ_4	ϖ_1

Moreover, for any representation $\rho : \mathfrak{g} \rightarrow \mathfrak{sl}(V)$, we have $d_{\rho} = 0 \bmod d_{\mathfrak{g}}$.

Proof. It is enough to calculate the Dynkin index for the fundamental weights (note that $d_{V \otimes W} = r_W d_V + r_V d_W$ if V and W are two $\mathfrak{g}$ -modules of rank r_V and r_W), which can be done explicitly [6]. We give the values here for E_8 , as not all of them in ([6], Table 5) are correct:

	d_{ϖ_1}	d_{ϖ_2}	d_{ϖ_3}	d_{ϖ_4}	d_{ϖ_5}	d_{ϖ_6}	d_{ϖ_7}	d_{ϖ_8}
E_8	1500	85500	5292000	8345660400	141605100	1778400	14700	60

3. The stack Bun_G .

Throughout this section k will be an algebraically closed field, G will be semi-simple algebraic group over k .

(3.1) Let X be a scheme over k . By a *principal G -bundle* over X (or just G -bundle for short), we understand a scheme $E \rightarrow X$ equipped with a right action of G such that, locally in the flat topology, E is trivial, *i.e.* isomorphic to $G \times X$ as a G -homogeneous space. In particular, E is affine, flat and smooth over X . Moreover, the above conditions imply that E is even locally trivial for the étale topology.

If F is a quasi-projective scheme on which G acts on the left and E is a G -bundle, we can form $E(F) = E \times^G F$ the *associated bundle with fiber F* . It is the quotient of $E \times F$ under the action of G defined by $g.(e, f) = (e.g, g^{-1}f)$. If H is a subgroup of G , the associated G/H -bundle $E(G/H)$ will be denoted simply by E/H .

Let $\rho : G \rightarrow G'$ be a morphism of algebraic groups. Then, as G acts on G' via ρ , we can form the *extension of the structure group* of a G -bundle E , that is the G' -bundle $E(G')$. Conversely, if F is a G' -bundle, a *reduction of structure group* F_G is a G -bundle E together with an isomorphism $F_G(G') \xrightarrow{\sim} F$. If ρ is a closed immersion, such reductions are in one to one correspondence with section of the associated bundle F/G .

(3.2) Let us collect some well known generalities on stacks for further reference. Let $\mathbb{A}ff/k$ be the flat affine site over k , that is the category of k -algebras equipped with the *fppf* topology. By k -space (resp. k -group) we understand a sheaf of sets (resp. groups) over $\mathbb{A}ff/k$. Any k -scheme can (and will) be considered as a k -space.

We will view k -stacks from the pseudo-functorial point of view, *i.e.* a k -stack $\mathfrak{X}$ will associate to every k -algebra R a groupoid $\mathfrak{X}(R)$ and to every morphism of k -algebras $u : R \rightarrow R'$ a functor $u^* : \mathfrak{X}(R') \rightarrow \mathfrak{X}(R)$ together with isomorphisms of functors $(u \circ v)^* \simeq v^* \circ u^*$ satisfying the usual cocycle condition. The required topological properties are that for every $x, y \in \text{ob } \mathfrak{X}(R)$ the presheaf $\underline{\text{Isom}}(x, y)$ is a sheaf and that all descent data are effective ([19], 2.1). Any k -space X may be seen as a k -stack, by considering a set as a groupoid (with the identity as the only morphism). Conversely, any k -stack $\mathfrak{X}$ such that $\mathfrak{X}(R)$ is a discrete groupoid (*i.e.* has only the identity as automorphisms) for all k -algebras R , is a k -space.

A morphism $F : \mathfrak{X} \rightarrow \mathfrak{Y}$ will associate, for every k -algebra R , a functor $\mathfrak{X}(R) \rightarrow \mathfrak{Y}(R)$ satisfying the obvious compatibility conditions. Let $S = \text{Spec}(R)$ and consider a morphism $\eta : S \rightarrow \mathfrak{Y}$, that is an object η of $\mathfrak{Y}(S)$. The fiber $\mathfrak{X}_\eta$ is a stack over S . The morphism F is *representable* if $\mathfrak{X}_\eta$ is representable as a scheme for all $S = \text{Spec}(R)$. A stack $\mathfrak{X}$ is *algebraic* if the diagonal morphism $\mathfrak{X} \rightarrow \mathfrak{X} \times \mathfrak{X}$ is representable, separated and quasi-compact and if there is a scheme X and a representable, smooth, surjective morphism of stacks $P : X \rightarrow \mathfrak{X}$.

Suppose X is a k -space and that the k -group Γ acts on X . Then the quotient stack $[X/\Gamma]$ is defined as follows. Let R be a k -algebra. The objects of $[X/\Gamma](R)$ are pairs (E, α) where E is a Γ -bundle over $\text{Spec}(R)$ and $\alpha : E \rightarrow X$ is G -equivariant, the arrows are defined in the obvious way and so are the functors $[X/\Gamma](R') \rightarrow [X/\Gamma](R)$.

(3.3) Let X be a smooth, complete and connected curve of genus g over k . We denote by Bun_G the stack of principal G -bundles over X which is defined as follows. For any k -algebra R denote X_R the scheme $X \times_k \text{Spec}(R)$. Then objects of $\text{Bun}_G(R)$ are G -bundles over X_R , morphisms of $\text{Bun}_G(R)$ are isomorphisms of G -bundles.

The following proposition is well known.

(3.4) PROPOSITION.— *The stack Bun_G is algebraic and smooth. Moreover we have $\dim \text{Bun}_G = (g - 1) \dim G$.*

(3.5) Choose a closed point p on X and set $X^* = X - p$. Let $\mathcal{O}$ be the completion of the local ring of X at p , and K its field of fractions. Set $D = \text{Spec}(\mathcal{O})$ and $D^* = \text{Spec}(K)$. We choose a local coordinate z at p and identify $\mathcal{O}$ with $k[[z]]$ and K with $k((z))$. Let R be a k -algebra. Define $X_R = X \times_k \text{Spec}(R)$, $X_R^* = X^* \times_k \text{Spec}(R)$, $D_R = \text{Spec}(R[[z]])$ and $D_R^* = \text{Spec}(R((z)))$. Then we have the cartesian diagram

$$\begin{array}{ccc}
D_R^* & \longrightarrow & D_R \\
\downarrow & & \downarrow \\
X_R^* & \longrightarrow & X_R
\end{array}$$

We denote by A_{X_R} the k -algebra $\Gamma(X_R^*, \mathcal{O}_{X_R^*})$.

(3.6) Loop groups. The category of k -spaces is closed under direct limits. A k -space (k -group) will be called an *ind-scheme* (resp. *ind-group*) if it is direct limit of a directed system of schemes. Remark that an ind-group is not necessarily an inductive limit of algebraic groups.

We denote by L_G the *loop group* of G that is the k -group defined $R \mapsto G(R((z)))$, where R is any k -algebra. The group of *positive loops*, that is the k -group $R \mapsto G(R[[z]])$ will be denoted by L_G^+ and the group of *negative loops*, that is the k -group $R \mapsto G(R[z^{-1}])$ will be denoted by L_G^- . The group of *loops coming from X^** , i.e. the k -group defined by $R \mapsto G(A_{X_R})$, will be denoted by L_G^X . Finally, we will use also the k -group L_G^{-} defined by $R \mapsto G(z^{-1}R[z^{-1}])$.

Choose a faithful representation $G \subset \mathrm{SL}_r$. For $N \geq 0$, we denote by $L_G(N)(R)$ the set of matrices $A(z)$ in $G(R((z))) \subset \mathrm{SL}_r(R((z)))$ such that for both $A(z)$ and $A(z)^{-1}$, the coefficients have a pole of order $\leq N$. This defines a subfunctor $L_G(N)$ of L_G which is obviously representable by an (infinite dimensional) affine k -scheme.

(3.7) PROPOSITION.— *The k -group L_G^+ is an affine group scheme. The k -group L_G is an ind-group, direct limit of the sequence of the schemes $(L_G(N))_{N \geq 0}$. Moreover, this ind-structure does not depend on the embedding $G \subset \mathrm{SL}_r$.*

The k -group L_G^X has the structure of an ind-group induced by the one of L_G . The quotient k -space $\mathcal{Q}_G := L_G/L_G^+$ has equally the structure of an ind-scheme: define $\mathcal{Q}_G(N) = L_G(N)/L_G^+$ (note that $L_G(N)$ is stable under right multiplication by $\mathcal{Q}_G(0) = L_G^+$).

(3.8) Consider triples (E, ρ, σ) where E is a G -bundle on X_R , $\rho : G \times X_R^* \rightarrow E|_{X_R^*}$ a trivialization of E over X_R^* and $\sigma : G \times D_R \rightarrow E|_{D_R}$ a trivialization of E over D_R . We let $T(R)$ be the set of isomorphism classes of triples (E, ρ, σ) .

(3.9) PROPOSITION.— *The ind-group L_G represents the functor T .*

Proof. Let (E, ρ, σ) be an element of $T(R)$. Pulling back the trivializations ρ and σ to D_R^* provides two trivializations ρ^* and σ^* of the pull back of E over D_R^* : these trivializations differ by an element $\gamma = \rho^{*-1} \circ \sigma^*$ of $G(R((z)))$.

Conversely, let us start from an element γ of $G(R((z)))$. This element defines an isomorphism of the pullbacks over D_R^* of the trivial G -bundle $\mathcal{F}$ over X_R^* and the trivial G -bundle $\mathcal{G}$

over D_R . These two torsors glue together to a G -bundle E in a functorial way by [4] (in fact [4] is written for SL_r but the extension to arbitrary G is straightforward).

These constructions are inverse to each other by construction. $\square$

(3.10) Consider the functor D_G which associates to a k -algebra R the set $D_G(R)$ of isomorphism classes of pairs (E, ρ) , where E is a G -bundle over X_R and ρ a trivialization of E over X_R^* .

(3.11) PROPOSITION.— *The ind-scheme $\mathcal{Q}_G$ represents the functor D_G .*

Proof. Let R be a k -algebra and q an element of $\mathcal{Q}_G(R)$. By definition there exists a faithfully flat homomorphism $R \rightarrow R'$ and an element γ of $G(R'((z)))$ such that the image of q in $\mathcal{Q}_G(R')$ is the class of γ . To γ corresponds by Proposition 3.9 a triple (E', ρ', σ') over $X_{R'}$. Let $R'' = R' \otimes_R R'$, and let $(E''_1, \rho''_1), (E''_2, \rho''_2)$ denote the pull-backs of (E', ρ') by the two projections of $X_{R''}$ onto $X_{R'}$. Since the two images of γ in $G(R''((z)))$ differ by an element of $G(R''[[z]])$, these pairs are isomorphic. So the isomorphism $\rho''_2 \rho''_1^{-1}$ over $X_{R''}^*$ extends to an isomorphism $u : E''_1 \rightarrow E''_2$ over $X_{R''}$, satisfying the usual cocycle condition (it is enough to check this over X^* , where it is obvious). Therefore (E', ρ') descends to a pair (E, ρ) on X_R as in the statement of the proposition.

Conversely, given a pair (E, ρ) as above over X_R , we can find a faithfully flat homomorphism $R \rightarrow R'$ and a trivialization σ' of the pull back of E over $D_{R'}$ (in fact, we can take R' to be the product of henselization of each localized ring R_x , $x \in \text{Spec}(R)$). By Proposition 3.9 we get an element γ' of $G(R'((z)))$ such that the two images of γ' in $G(R''((z)))$ (with $R'' = R' \otimes_R R'$) differ by an element of $G(R''[[z]])$; this gives an element of $\mathcal{Q}_G(R)$. The two constructions are clearly inverse one of each other. $\square$

We will make use of the following theorem

(3.12) THEOREM.— *(Drinfeld-Simpson [8]) Let E be a G -bundle over X_R . Then the restriction of E to X_R^* is trivial fppf locally over $\text{Spec}(R)$. If $\text{char}(k)$ does not divide the order of $\pi_1(G(\mathbb{C}))$, then this is even true étale locally.*

(3.13) Proof of Theorem 1.3. The universal G -bundle E over $X \times \mathcal{Q}_G$ (Proposition 3.11), gives rise to a map $\pi : \mathcal{Q}_G \rightarrow \text{Bun}_G$. This map is L_G^X -invariant, hence induces a morphism of stacks $\bar{\pi} : L_G^X \backslash \mathcal{Q}_G \rightarrow \text{Bun}_G$.

On the other hand we can define a map $\text{Bun}_G \rightarrow L_G^X \backslash \mathcal{Q}_G$ as follows. Let R be a k -algebra, E a G -bundle over X_R . For any R -algebra R' , let $T(R')$ be the set of trivializations ρ of $E_{R'}$ over $X_{R'}^*$. This defines a R -space T on which the group L_G^X acts. By Theorem 3.12, it is a

torsor under L_G^X . To any element of $T(R')$ corresponds a pair $(E_{R'}, \rho)$, hence by Proposition 3.11 an element of $\mathcal{Q}_G(R')$. In this way we associate functorially to an object E of $\text{Bun}_G(R)$ a L_G^X -equivariant map $\alpha : T \rightarrow \mathcal{Q}_G$. This defines a morphism of stacks $\text{Bun}_G \rightarrow L_G^X \backslash \mathcal{Q}_G$ which is the inverse of $\bar{\pi}$.

The second assertion means that for any scheme S over k (resp. over k such that $\text{char}(k)$ does not divide the order of $\pi_1(G(\mathbb{C}))$) and any morphism $f : T \rightarrow \text{Bun}_G$, the pull back to S of the fibration π is *fppf* (resp. *étale*) locally trivial, i.e. admits local sections (for the *fppf* (resp. *étale*) topology). Now f corresponds to a G -bundle E over $X \times S$. Let $s \in S$. Again by Theorem 3.12, we can find an *fppf* (resp. *étale*) neighborhood U of s in S and a trivialization ρ of $E|_{X \times U}$. The pair (E, ρ) defines a morphism $g : U \rightarrow \mathcal{Q}_G$ (Proposition 3.11) such that $\pi \circ g = f$, that is a section over U of the pull back of the fibration π . $\square$

4. The infinite grassmannian $\mathcal{Q}_G$

Let G be semi-simple and k be an algebraically closed field of characteristic 0 in (4.2).

(4.1) We will use the following two facts:

(a) We may write $\mathcal{Q}_G$ as direct limit of projective finite-dimensional k -schemes.

(b) The multiplication map $\mu : L_G^{-} \times L_G^{+} \rightarrow L_G$ is an open immersion.

For the first statement, remark that it is enough to consider the k -space $\mathcal{Q}(\mathfrak{g})$ parameterizing isomorphism classes of sheaves of Lie algebra which are locally of the form $S \times \text{Lie}(G)$ together with a trivialization over $\mathbb{P}_1^*$ (note that, looking at the adjoint group, $\mathcal{Q}_G$ may be seen as a connected component of $\mathcal{Q}(\mathfrak{g})$) then argue as in [3]. For the second statement, the argument of ([3] Proposition 1.11) generalizes to arbitrary G , once we know the following.

Claim. Suppose Y is a proper S -scheme and that the structural morphism has a section $\sigma : S \rightarrow Y$. Suppose moreover that $G \backslash H$ is a reductive subgroup of a reductive group H . Then, for any G -bundle P trivial along σ the following is true: if the associated H -bundle $P(H)$ is trivial, the G -bundle P is so.

Indeed, by assumption, there exists a section $\tau : Y \rightarrow P(H)$. The quotient $G \backslash H$ is affine. Therefore, the composite morphism from Y to $G \backslash H$ (which is the composition of τ and of the canonical projection $P(H) \rightarrow G \backslash H$) factors as $Y \rightarrow S \xrightarrow{p} G \backslash H$. After an eventual translation of τ by an element of $H(S)$, we can assume that the restriction $\sigma^*(\tau)$ of τ along σ is induced by the trivialization of P . Therefore, the morphism p is the constant morphism with constant value $G \in G \backslash H$. In other words, locally for the étale topology on Y , the section τ can be written

$$\tau(y) = (p(y), h(y)) \text{ mod } G \text{ where } y \in Y \text{ and } p(y) \in P, h(y) \in G.$$

The expression $p(y)h(y)^{-1}$ is well defined and defines a section of P .

(4.2) The quotient L_G/L_G^+ has also been studied by Kumar and Mathieu. But the structure of ind-variety they put on the quotient is, a priori, not the same as the functorial one of section 3. As we will use their results, we have to identify them.

PROPOSITION.— *The ind-structure of $\mathcal{Q}_G$ defined in section 3 coincides with the ind-structure of Kumar and Mathieu.*

Proof. Recall that an ind-scheme is called *reduced* (resp. *irreducible*, *integral*) if it is a direct limit of an increasing sequence of reduced (resp. irreducible, integral) schemes. By Lemma 6.3 of [3] an ind-scheme is integral if and only if it is irreducible and reduced. According to Faltings [10] (see [3] for the case SL_r), the ind-group L_G^- is integral. This may be seen by looking at $(L_G^-)_{red}$ and using Shavarevich's theorem that a closed immersion of irreducible ind-affine groups which is an isomorphism on Lie algebras, is an isomorphism [27]. Note that irreducibility is due to the fact that any element can be deformed to a constant in G ; that $\text{Lie}(L_G^-) \rightarrow \text{Lie}(L_G^-)_{red}$ is an isomorphism can be seen by using the fact that $\text{Lie}(G)$ is generated by nilpotent elements.

It follows for semisimple G that L_G^{-} , which is a semidirect product of G and L_G^- , is integral, and furthermore if G is simply connected then $\mathcal{Q}_G$ is integral. Indeed by (4.1 b) $L_G^{-} \rightarrow \mathcal{Q}_G$ is an open immersion hence it is enough to show that $\mathcal{Q}_G$ is irreducible. Using that connected ind-groups are irreducible ([27], Proposition 3) and the quotient morphism $L_G \rightarrow \mathcal{Q}_G$ we reduce to prove the connectedness of L_G which is well known (and follows for example from uniformization for $\mathbb{P}_1$ and the corresponding statement for $\text{Bun}_G(\mathbb{P}_k^1)$, cf. [8]).

We are ready to deduce the identification of our ind-structure on $\mathcal{Q}_G$ with the one used by Kumar or Mathieu in their generalized Borel-Weil theory. Both Kumar and Mathieu define the structure of ind-variety on L_G/L_G^+ using representation theory of Kac-Moody algebras; for instance Kumar, following Slodowy [28], considers the basic representation $\mathcal{H}_\ell$ for a fixed ℓ , and a highest weight vector v_ℓ . The subgroup L_G^+ is the stabilizer of the line kv_ℓ in $\mathbb{P}(\mathcal{H}_\ell)$, so the map $g \mapsto gv_\ell$ induces an injection $i_\ell : L_G/L_G^+ \hookrightarrow \mathbb{P}(\mathcal{H}_\ell)$. Let U be the subgroup of L_G^+ consisting of elements $A(z)$ such that $A(0)$ is in the unipotent part of a fixed Borel subgroup $B \subset G$; to each element w of the Weyl group is associated a ‘‘Schubert variety’’ X_w which is a finite union of orbits of U . It turns out that the image under i_ℓ of X_w is actually contained in some finite-dimensional projective subspace $\mathbb{P}_w$ of $\mathbb{P}(\mathcal{H}_\ell)$, and is Zariski closed in $\mathbb{P}_w$. This defines on X_w a structure of reduced projective variety, and a structure of ind-variety on $L_G/L_G^+ = \varinjlim X_w$.

By a result due to Faltings (cf. the Appendix of [3] for SL_r), the irreducible integrable representation $\mathcal{H}_\ell$ of $\widehat{L_{\mathfrak{g}}}$ can be ‘‘integrated’’ to an *algebraic* projective representation of L_G , that is a morphism of k -groups $L_G \rightarrow \text{PGL}(\mathcal{H}_\ell)$. It follows that the map i_c is a morphism of ind-schemes of $\mathcal{Q}_G$ into $\mathbb{P}(\mathcal{H}_\ell)$ (which is the direct limit of its finite-dimensional subspaces).

But i_ℓ is even an *embedding*. It is injective by what we said above; let us check that it induces an injective map on the tangent spaces. Since it is equivariant under the action of L_G it is enough to prove this at the origin ω of $\mathcal{Q}_G$. Then it follows from the fact that the annihilator of v_ℓ in the Lie algebra $L_{\mathfrak{g}}$ is $L_{\mathfrak{g}}^+$.

Therefore the restriction of i_ℓ to each of the subvarieties $\mathcal{Q}_G(N)$ is proper, injective, and injective on the tangent spaces, hence is an embedding (in some finite-dimensional projective subspace of $\mathbb{P}(\mathcal{H}_\ell)$). Each X_w is contained in some $\mathcal{Q}_G(N)$, and therefore is a closed subvariety of $\mathcal{Q}_G(N)_{\text{red}}$. Each orbit of U is contained in some X_w ; since the X_w 's define an ind-structure, each $\mathcal{Q}_G(N)$ is contained in some X_w , so that $\mathcal{Q}_G(N)_{\text{red}}$ is a subvariety of X_w . Since $\mathcal{Q}_G$ is the direct limit of the $\mathcal{Q}_G(N)_{\text{red}}$, the two ind-structures coincide. $\square$

5. The ind-group L_G^X

Throughout this section we suppose $k = \mathbb{C}$ and G semi-simple and simply connected.

(5.1) PROPOSITION.— *The ind-group L_G^X is integral.*

(5.2) COROLLARY.— *Every character $\chi : L_G^X \rightarrow G_m$ is trivial.*

Proof. The differential of χ , considered as a function on L_G^X , is everywhere vanishing. Indeed, since χ is a group morphism, this means that the deduced Lie algebra morphism $\mathfrak{g} \otimes A_X \rightarrow k$ is zero. But as the derived algebra $\mathcal{D}(\mathfrak{g} \otimes A_X)$ is $\mathcal{D}(\mathfrak{g}) \otimes A_X$ and therefore equal to $\mathfrak{g} \otimes A_X$ because $\mathfrak{g}$ is simple, any Lie algebra morphism $\mathfrak{g} \otimes A_X \rightarrow k$ is trivial.

As L_G^X is integral we can write L_G^X as the direct limit of integral varieties V_n . The restriction of χ to V_n has again zero derivative and is therefore constant. For large n , the varieties V_n contain 1. This implies $\chi|_{V_n} = 1$ and we are done. $\square$

Proof. To see that the ind-group L_G^X is reduced, consider the étale trivial morphism $\bar{\pi} : \mathcal{Q} \rightarrow \text{Bun}_G$. Locally for the étale topology, $\bar{\pi}$ is a product $\Omega \times L_G^X$ (where Ω is an étale neighborhood of Bun_G). Then use that $\mathcal{Q}$ is reduced by (4.2). As connected ind-groups are irreducible by Proposition 3 of [27] it is enough to show that L_G^X is connected.

The idea how to prove that L_G^X is connected is due to V. Drinfeld: consider distinct points $p_1, \dots, p_i$ of X which are all distinct from p . Define $X_i^* = X - \{p, p_1, \dots, p_i\}$ and, for every k -algebra R , define $X_{i,R}^* = X_i^* \times_k \text{Spec}(R)$. Denote by $A_{X_{i,R}}$ the k -algebra $\Gamma(X_{i,R}^*, \mathcal{O}_{X_{i,R}^*})$ and by $L_G^{X,i}$ the k -group $R \mapsto G(A_{X_{i,R}})$. As L_G^X , the k -group $L_G^{X,i}$ is an ind-group (cf. 3.6). The natural inclusion $A_{X_{i,R}} \subset A_{X_{i+1,R}}$ defines a closed immersion $f : L_G^{X,i} \rightarrow L_G^{X,i+1}$.

(5.3) LEMMA.— *The closed immersion $L_G^{X,i} \rightarrow L_G^{X,i+1}$ defines a bijection*

$$\pi_0(L_G^{X,i}) \xrightarrow{\sim} \pi_0(L_G^{X,i+1}).$$

Proof. Consider the morphism $L_G^{X,i+1} \rightarrow L_G$ defined by the developpement in Laurent series at p_{i+1} . We get a morphism $\phi_{i+1} : L_G^{X,i+1} \rightarrow \mathcal{Q}_G^{p_{i+1}}$, where we denote $\mathcal{Q}_G^{p_{i+1}} = L_G/L_G^+$. (of course $\mathcal{Q}_G = \mathcal{Q}_G^{p_{i+1}}$ but we emphasize here that we will consider the point p_{i+1} and not p .)

Claim: The morphism $\phi_{i+1} : L_G^{X,i+1} \rightarrow \mathcal{Q}_G^{p_{i+1}}$ induces an isomorphism on the level of stacks $\bar{\phi}_{i+1} : L_G^{X,i+1}/L_G^{X,i} \simeq \mathcal{Q}_G^{p_{i+1}}$ and is locally trivial for the étale topology.

The lemma reduces to the claim. Indeed, as G is semi-simple and simply connected, we have $\pi_i([\mathcal{Q}_G^{p_{i+1}}]^{an}) = 1$ for $i = 0, 1$ (by (4.2) and Kumar and Mathieu) and the exact homotopy sequence associated of the (Serre)-fibration ϕ_{i+1} shows that $\pi_0([L_G^{X,i}]^{an}) \xrightarrow{\sim} \pi_0([L_G^{X,i+1}]^{an})$ ($[]^{an}$ means we consider the usual topology). From the bijection $\pi_0(L_G^{X,i}(N)) \xrightarrow{\sim} \pi_0([L_G^{X,i}(N)]^{an})$ and Proposition 2 of [27] it follows then that $\pi_0(L_G^{X,i}) \rightarrow \pi_0([L_G^{X,i}]^{an})$ is bijective.

Proof of the claim: Clearly $\phi_{i+1} : L_G^{X,i+1} \rightarrow \mathcal{Q}_G^{p_{i+1}}$ is $L_G^{X,i}$ invariant, hence defines a map $\bar{\phi}_{i+1} : L_G^{X,i+1}/L_G^{X,i} \rightarrow \mathcal{Q}_G^{p_{i+1}}$. Define a morphism $\mathcal{Q}_G^{p_{i+1}} \rightarrow L_G^{X,i+1}/L_G^{X,i}$ as follows. Let R be a k -algebra. By Proposition 3.11 to an element of $\mathcal{Q}_G^{p_{i+1}}(R)$ corresponds a G -bundle $E \rightarrow X_R$ together with a trivialization $\tau_{p_{i+1}} : G \times X_{p_{i+1},R}^* \rightarrow E|_{X_{p_{i+1},R}^*}$. Here by $X_{p_{i+1},R}^*$ we denote $(X - \{p_{i+1}\}) \times_k \text{Spec}(R)$. For any R -algebra R' , denote $T(R')$ the set of trivializations τ_i of $E_{R'}$ over $X_{i,R}^*$. This defines a R -space T on which $L_G^{X,i}$ acts. By Theorem 3.12 it is a torsor under $L_G^{X,i}$. For any $\tau_i \in T(R')$ the composite $\tau_i^{-1} \circ \tau_{p_{i+1}}$ defines a morphism $X_{i+1,R}^* \rightarrow G$ hence an element of $L_G^{X,i+1}(R)$. In this way we associate functorially to an object $(E, \tau_{p_{i+1}})$ of $\mathcal{Q}_G^{p_{i+1}}(R)$ a $L_G^{X,i}$ -invariant map $\alpha : T \rightarrow L_G^{X,i+1}$, which defines the inverse of $\bar{\phi}$. The assertion concerning the local triviality is proved as in Theorem 1.3. $\square$

Let us show that every element $g \in L_G^X(k)$ is in the connected component of the unit of $L_G^X(k)$. Using the well known fact that $G(K)$ is generated by the standard unipotent subgroups $U_\alpha(K)$, $\alpha \in \Delta$, we may suppose that g is of the form $\prod_{j \in J} \exp(f_j n_j)$ where the n_j are nilpotent elements of $\mathfrak{g}$ and $f_j \in K$. Let $\{p_1, \dots, p_i\}$ be the poles of the functions $f_j, j \in J$. The morphism

$$\begin{array}{ccc} \mathbb{A}^1 & \longrightarrow & L_G^{X,i} \\ t & \longmapsto & \prod_{j \in J} \exp(t f_j n_j) \end{array}$$

is a path from g to 1 in $L_G^{X,i}$. By Lemma 5.3, the morphism $\pi_0(L_G^X) \rightarrow \pi_0(L_G^{X,i})$ is bijective which proves that g and 1 are indeed in the same connected component of L_G^X . $\square$

6. Pfaffians

Let k be an algebraically closed field of characteristic $\neq 2$ and S a k -scheme.

(6.1) The Picard categories

Let A be $\mathbb{Z}$ or $\mathbb{Z}/2\mathbb{Z}$. Denote by $\mathfrak{L}_A$ the groupoid of A -graded invertible $\mathcal{O}_S$ -modules. The objects of $\mathfrak{L}_A$ are pairs $[L] = (L, a)$ of invertible $\mathcal{O}_S$ -modules L and locally constant functions $a : S \rightarrow A$, morphisms $[f] : [L] \rightarrow [M]$ are defined if $a = b$ and are isomorphisms $f : L \rightarrow M$ of $\mathcal{O}_S$ -modules. Denote $\mathbb{I}_A$ the object $(\mathcal{O}_S, 0)$. The category $\mathfrak{L}_A$ has tensor products, defined by $[L] \otimes [M] = (L \otimes M, a + b)$. Given $[L]$ and $[M]$ we have Koszul's symmetry isomorphism $\sigma_{[L],[M]} : [L] \otimes [M] \rightarrow [M] \otimes [L]$ defined on local sections ℓ and m by $\sigma_{L,M}(\ell \otimes m) = (-1)^{ab} m \otimes \ell$.

Denote $\det_A$ the functor from the category of coherent locally free $\mathcal{O}_S$ -modules with isomorphisms defined by $\det_A = (\Lambda^{max}, \text{rang}(V))$ and $\det_A(f) = \Lambda^{max}(f)$.

In the following we drop the subscript A for $A = \mathbb{Z}$ and replace it by 2 for $A = \mathbb{Z}/2\mathbb{Z}$.

(6.2) Pfaffians

Let V be a coherent locally free $\mathcal{O}_S$ -module of rank $2n$. Let $\text{Pf} : \Lambda^2 V^* \rightarrow \Lambda^{2n} V^*$ be the unique map that commutes with base changes and such that if $(e_1^*, \dots, e_{2n}^*)$ is a local frame of V^* and $\alpha = \sum_{i < j} a_{ij} e_i^* \wedge e_j^*$, then

$$\text{Pf}(\alpha) = \text{pf}(a) e_1^* \wedge \dots \wedge e_{2n}^*$$

where $\text{pf}(a)$ is the pfaffian [Bourbaki, Algèbre 9.5.2] of the alternating matrix $a_{ij} = -a_{ji}$ for $i > j$ and $a_{ii} = 0$ for $i = 1, \dots, 2n$.

Suppose $\alpha : V \rightarrow V^*$ is skewsymmetric. View α as a section of $\Lambda^2 V^*$ and define the *pfaffian* of α as the section $\text{Pf}(\alpha) : \mathcal{O}_S \rightarrow \Lambda^{2n} V^*$. By [Bourbaki, Algèbre 9.5.2] we know that

$$\begin{array}{ccc} \mathcal{O}_S \otimes \mathcal{O}_S & \longrightarrow & \mathcal{O}_S \\ \text{Pf}(\alpha) \otimes \text{Pf}(\alpha) \downarrow & & \swarrow \det(\alpha) \\ \Lambda^{2n} V^* \otimes \Lambda^{2n} V^* & & \end{array} \quad (6.2.1)$$

commutes and that, if u is an endomorphism of V^* , then

$$\begin{array}{ccc} \mathcal{O}_S & \xrightarrow{\text{Pf}(\alpha)} & \Lambda^{2n} V^* \\ \text{Pf}(u\alpha u^*) \downarrow & & \swarrow \det(u) \\ \Lambda^{2n} V^* & & \end{array} \quad (6.2.2)$$

commutes.

(6.3) *The pfaffian functor*

We consider the following category $\mathcal{A} = \mathcal{A}^\bullet(S)$: objects are complexes of locally free coherent $\mathcal{O}_S$ -modules concentrated in degrees 0 and 1 of the form

$$0 \longrightarrow E \xrightarrow{\alpha} E^* \longrightarrow 0$$

with α skewsymmetric. Morphisms between two such complexes $E^\bullet$ and $F^\bullet$ are morphisms of complexes $f^\bullet : E^\bullet \longrightarrow F^\bullet$ such that $f^{\bullet*}[-1]$ is a homotopy inverse of $f^\bullet$, i.e. $f^{\bullet*}[-1] \circ f^\bullet$ and $f^\bullet \circ f^{\bullet*}[-1]$ are homotopic to the identity.

Let $\pi : \mathfrak{L} \rightarrow \mathfrak{L}_2$ be the projection functor, $\Delta : \mathfrak{L}_2 \rightarrow \mathfrak{L}_2$ be the functor defined by $[L] \mapsto [L] \otimes [L]$ and $[f] \mapsto [f] \otimes [f]$ and $\text{Det} : \mathcal{A} \rightarrow \mathfrak{L}$ be the determinant functor [15].

(6.4) PROPOSITION.— *There is a natural functor, $\text{Pf} : \mathcal{A}^\bullet \rightarrow \mathfrak{L}_2$, commuting with base changes, and a natural isomorphism of functors:*

$$\pi \circ \text{Det} \xrightarrow{\sim} \Delta \circ \text{Pf}.$$

Moreover, if $f^\bullet : E^\bullet \longrightarrow F^\bullet$ is homotopic to the identity then $\text{Pf}(f^\bullet) = \text{id}$.

Proof. Define Pf on the level of objects by $\text{Pf}(E^\bullet) = \det_2(E)$. On the level of morphisms we do the following.

Let $f^\bullet = (f_0, f_1) : E^\bullet \longrightarrow F^\bullet$ be a morphism of $\mathcal{A}^\bullet$:

$$\begin{array}{ccc} E & \xrightarrow{\alpha_E} & E^* \\ f_0 \downarrow & & f_1 \downarrow \\ F & \xrightarrow{\alpha_F} & F^* \end{array}$$

and consider the complex $C_f^\bullet$ (which is up to sign the cone of $f^\bullet$)

$$C_f^\bullet = 0 \longrightarrow E \xrightarrow{\begin{pmatrix} \alpha_E \\ -f_0 \end{pmatrix}} E^* \oplus F \xrightarrow{(f_1 \ \alpha_F)} F^* \longrightarrow 0$$

As $f^\bullet$ is a quasi-isomorphism, $C_f^\bullet$ is acyclic. By the usual additivity property of determinants, we get a canonical isomorphism

$$d(f) : \Lambda^{\max} E \otimes \Lambda^{\max} F^* \rightarrow \Lambda^{\max} E^* \otimes \Lambda^{\max} F.$$

Recall that this isomorphism is defined by taking a section $\begin{pmatrix} u \\ v \end{pmatrix}$ of $(f_1 \ \alpha_F)$ and calculating the determinant, which is independent of this choice, of the morphism

$$M(f) = \begin{pmatrix} \alpha & u \\ -f_0 & v \end{pmatrix} \in \text{Hom}(E \oplus F^*, E^* \oplus F)$$

(6.5) LEMMA.— *There is a skew-symmetric morphism $\gamma_f \in \text{Hom}(F^*, F)$ such that $\begin{pmatrix} f_0^* \\ \gamma \end{pmatrix}$ is a section of $(f_1 \alpha_F)$.*

Proof. As $f \circ f^*[-1]$ is homotopic to Id there is a morphism h such that $f_0 f_1^* - 1 = h \alpha_F$ and $f_1 f_0^* - 1 = \alpha_F h$. Now define $\gamma_f = \frac{h^* - h}{2}$. $\square$

The pfaffian of the skew-symmetric morphism

$$M(f, \gamma_f) = \begin{pmatrix} \alpha & f_0^* \\ -f_0 & \gamma_f \end{pmatrix} \in \text{Hom}(E \oplus F^*, E^* \oplus F)$$

defines a section $\text{pf}(M(f, \gamma_f)) : \mathcal{O}_S \rightarrow \Lambda^{\max} E^* \otimes \Lambda^{\max} F$.

(6.6) LEMMA.— *The section $\text{pf}(M(f, \gamma_f))$ is independent of the choice of γ_f .*

Proof. Suppose γ'_f is another morphism satisfying (6.5). Then there is $g \in \text{Hom}(F^*, E)$ such that $\alpha_E g = 0$ and $f_0 g = -g^* f_0^*$ [use that γ_f and γ'_f are skew]. These relations give

$$M(f, \gamma'_f) = \begin{pmatrix} 1 & 0 \\ \frac{g^*}{2} & 1 \end{pmatrix} M(f, \gamma_f) \begin{pmatrix} 1 & \frac{g}{2} \\ 0 & 1 \end{pmatrix}$$

which in turn implies the required equality by (6.2.2). $\square$

As $\text{rank}(E) = \text{rank}(F) \bmod 2$, we get the isomorphism in $\mathfrak{L}_2$:

$$\text{pf}(M(f)) : \mathbb{I}_2 \xrightarrow{\sim} \det_2(E)^* \otimes \det_2(F).$$

Define the pfaffian of $f^\bullet$ by

$$\text{Pf}(f^\bullet) : \det_2(E) \xrightarrow{1 \otimes \text{pf}(M(f, \gamma_f))} \det_2(E) \otimes \det_2(E)^* \otimes \det_2(F) \xrightarrow{\text{ev}_{\det_2(E)}} \det_2(F)$$

(6.7) LEMMA.— *$\text{Pf} : \mathcal{A} \rightarrow \mathfrak{L}_2$ defines a functor.*

Proof. As $\text{pf}(M(\text{Id}, 0)) = 1$, we have $\text{Pf}(\text{Id}) = \text{Id}$. Let $f^\bullet : E^\bullet \rightarrow F^\bullet$ and $g : F^\bullet \rightarrow G^\bullet$ be two morphisms of $\mathcal{A}$. Then the following diagram is commutative

$$\begin{array}{ccc} \mathbb{I}_2 & \xrightarrow{\text{pf}(M(f, \gamma_f)) \otimes \text{pf}(M(g, \gamma_g))} & \det_2(E)^* \otimes \det_2(F) \otimes \det_2(F)^* \otimes \det_2(G) \\ \text{Id} \downarrow & & \downarrow 1 \otimes \text{ev}_{\det_2(F)} \otimes 1 \\ \mathbb{I}_2 & \xrightarrow{\text{pf}(M(g \circ f, \gamma_{g \circ f}))} & \det_2(E)^* \otimes \det_2(G) \end{array}$$

Indeed, remark that $\gamma_{gf} = g_0\gamma_f g_0^* + \gamma_g$ satisfies (6.5) for $g \circ f$ and make use of (6.2.2) first with

$$\begin{pmatrix} \alpha_E & f_0^* & 0 & f_0^* g_0^* \\ -f_0 & \gamma_f & 1 & \gamma_f g_0^* \\ 0 & -1 & 0 & 0 \\ -g_0 f_0 & g_0 \gamma_f & 0 & \gamma_{gf} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -f_1 & -\alpha_F & 1 & 0 \\ 0 & g_0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha_E & f_0^* & 0 & 0 \\ -f_0 & \gamma_f & 0 & 0 \\ 0 & 0 & \alpha_F & g_0^* \\ 0 & 0 & -g_0 & \gamma_g \end{pmatrix} \begin{pmatrix} 1 & 0 & -f_1^* & 0 \\ 0 & 1 & \alpha_F & g_0^* \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

and then with

$$\begin{pmatrix} \alpha_E & 0 & 0 & f_0^* g_0^* \\ 0 & \gamma_f & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -g_0 f_0 & 0 & 0 & \gamma_{gf} \end{pmatrix} = \begin{pmatrix} 1 & 0 & f_0^* & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & g_0 \gamma_f & 1 \end{pmatrix} \begin{pmatrix} \alpha_E & f_0^* & 0 & f_0^* g_0^* \\ -f_0 & \gamma_f & 1 & \gamma_f g_0^* \\ 0 & -1 & 0 & 0 \\ -g_0 f_0 & g_0 \gamma_f & 0 & \gamma_{gf} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ f_0 & 0 & 1 & -\gamma_f g_0^* \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

The commutativity of the above diagram shows $\text{Pf}(g \circ f) = \text{Pf}(g) \circ \text{Pf}(f)$. $\square$

The statement on the natural transformation follows from the definitions and (6.2.1). It remains to prove that if $f^\bullet : E^\bullet \rightarrow E^\bullet$ is homotopic to the identity, then $\text{Pf}(f) = \text{Id}$. Indeed, let $h : E^* \rightarrow E$ be such that $f_0 - h\alpha_E = 1$ and $f_1 - \alpha h = 1$. Then $\gamma_f := -h + f_0 h^*$ satisfies (6.5) and the statement follows from (6.2.2) and

$$\begin{pmatrix} a_E & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ h & 1 \end{pmatrix} \begin{pmatrix} a_E & f_0^* \\ -f_0^* & \gamma_f \end{pmatrix} \begin{pmatrix} 1 & h^* \\ 0 & 1 \end{pmatrix}$$

This completes the proof of Proposition 6.4. $\square$

7. Line bundles on Bun_G .

Let k be an algebraically closed field of characteristic $\neq 2$ in Sections (7.2)-(7.6). Denote $\text{Pic}(\text{Bun}_G)$ the group of isomorphism classes of line bundles on Bun_G . (See ([3], 3.7) for a discussion of line bundles over k -spaces and stacks). We will construct special elements of $\text{Pic}(\text{Bun}_G)$.

(7.1) The determinant line bundle.

We start with the well known case of $G = \text{GL}_r$: let $\mathcal{F}$ be a family of vector bundle of rank r parameterized by the locally noetherian k -scheme S . Recall that the complex $\text{Rpr}_{1*}(\mathcal{F})$ may be represented by a perfect complex of length one $K^\bullet$ and define $\mathcal{D}_{\mathcal{F}}$ to be $\det(K^\bullet)^{-1}$. This does not, up to canonical isomorphism, depend on the choice of $K^\bullet$. As the formation of the determinant commutes with base change, the fiber of $\mathcal{D}_{\mathcal{F}}$ over the point $s \in S$ is $\Lambda^{\max} H^0(X, \mathcal{F}(s))^* \otimes \Lambda^{\max} H^1(X, \mathcal{F}(s))$. The line bundle $\mathcal{D}_{\mathcal{F}}$ is called the *determinant of cohomology* line bundle associated to the family $\mathcal{F}$.

Let $\mathcal{U}$ be the universal vector bundle on $\text{Bun}_{\text{GL}_r} \times X$ and define the determinant line bundle $\mathcal{D} = \det(\text{Rpr}_{1*}\mathcal{U})^{-1}$. It has the following universal property: for every family $\mathcal{F}$ of vector bundle parameterized by the locally noetherian k -scheme S , we have $f_{\mathcal{F}}^*(\mathcal{D}) = \mathcal{D}_{\mathcal{F}}$ where $f_{\mathcal{F}} : S \rightarrow \text{Bun}_{\text{GL}_r}$ is the deduced modular morphism.

For the case of general G , consider a representation $\rho : G \rightarrow \text{GL}_r$ and consider the morphism obtained by extension of structure group $f_{\rho} : \text{Bun}_G \rightarrow \text{Bun}_{\text{GL}_r}$. Then define the determinant of cohomology associated to ρ by $\mathcal{D}_{\rho} = f_{\rho}^*(\mathcal{D})$.

(7.2) The Pfaffian bundle

Consider $G = \text{Spin}_r$ with $r \geq 3$ (resp. $G = G_2$). Then the standard representation ϖ_1 factors through SO_r (resp. SO_7). The stack Bun_{SO_r} has two components: $\text{Bun}_{\text{SO}_r}(0)$ and $\text{Bun}_{\text{SO}_r}(1)$. They are distinguished by the second Stiefel-Whitney class

$$w_2 : H_{\text{ét}}^1(X, \text{SO}_r) \rightarrow H_{\text{ét}}^2(X, \mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}.$$

Let κ be a theta-characteristic on X . Twisting by κ , we may and will see a SO_r -bundle as a vector bundle F with trivial determinant together with a *symmetric* isomorphism $\sigma : F \rightarrow F^{\vee}$, where $F^{\vee} = \underline{\text{Hom}}_{\mathcal{O}_X}(F, \omega_X)$. The following Proposition shows the existence, for every κ , of a canonical square root $\mathcal{P}_{\kappa}$ of the determinant bundle $\mathcal{D}_{\varpi_1}$ over Bun_{SO_r} .

(7.3) PROPOSITION.— *Let (F, σ) be a family of vector bundles F equipped with a quadratic form σ with values in ω_X parameterized by the locally noetherian k -scheme S . Then the determinant of cohomology $\mathcal{D}_F$ admits a canonical square root $\mathcal{P}_{(F, \sigma)}$. Moreover, if $f : S' \rightarrow S$ is a morphism of locally noetherian k -schemes then we have $\mathcal{P}_{(f^*F, f^*\sigma)} = f^*\mathcal{P}_{(F, \sigma)}$.*

Proof. By ([30], prop. 2.1 and proof of corollary 2.2, cf. also [14]), Zariski locally on S , there are length 1 complexes $M^{\bullet}$ of finite free $\mathcal{O}_S$ -modules and quasi-isomorphisms $f : M^{\bullet} \rightarrow \text{Rpr}_{1*}(F)$ such that the composition in the derived category $D(S)$ (use σ and Grothendieck duality)

$$M^{\bullet} \xrightarrow{f} \text{Rpr}_{1*}(F) \xrightarrow{\tau} \text{R}\underline{\text{Hom}}^{\bullet}(\text{Rpr}_{1*}(F), \mathcal{O}_S)[-1] \xrightarrow{f^{*}[-1]} M^{\bullet*}[-1]$$

lifts to a symmetric isomorphism of complexes $\varphi : M^{\bullet} \rightarrow M^{\bullet*}[-1]$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & M^0 & \xrightarrow{d_{M^{\bullet}}} & M^1 & \longrightarrow & 0 \\ & & \varphi_0 \downarrow \wr & \searrow \alpha & \wr \downarrow \varphi_0^* & & \\ 0 & \longrightarrow & M^{1*} & \xrightarrow{-d_{M^{\bullet*}}} & M^{0*} & \longrightarrow & 0 \end{array}$$

Define $\widetilde{M}^{\bullet}$ by $0 \rightarrow M^0 \xrightarrow{\alpha} M^{0*} \rightarrow 0$. Then α is skew and we have a natural isomorphism of complexes $\psi : M^{\bullet} \rightarrow \widetilde{M}^{\bullet}$ such that $\psi^*[-1]\psi = f^*[-1]\tau f$ in $D^b(S)$.

Cover S by open subsets U_i together with complexes $(M_i^\bullet, d_{M_i^\bullet})$ and quasi-isomorphisms $f_i : M_i^\bullet \rightarrow Rpr_{1*}(F)|_{U_i}$ as above. We define $\mathcal{P}_{(F,\sigma)}$ over U_i by $\mathcal{P}_{i,(F,\sigma)} = \text{Pf}(\widetilde{M}_i^\bullet)$ and construct patching data $\rho_{ij} : \mathcal{P}_{i,(F,\sigma)} \xrightarrow{\sim} \mathcal{P}_{j,(F,\sigma)}$ over $U_{ij} = U_i \cap U_j$ in the following way. Define first the morphism of complexes $\Sigma_{ij} : \widetilde{M}_i \rightarrow \widetilde{M}_j$ as a lifting of the isomorphism in $D^b(U_{ij})$

$$\psi_j^{*-1}[-1]f_j^*[-1]\tau f_i\psi_i^{-1},$$

then ρ_{ij} by $\text{Pf}(\Sigma_{ij})$ [note that it follows from the symmetry of σ (and that the components of the $\widetilde{M}_i$ are free) that Σ_{ij} is a morphism of $\mathcal{A}(U_{ij})$]. By 6.4, ρ_{ij} does not depend on the particular chosen lifting and the functoriality of Pf translates into $\rho_{ii} = \text{Id}$, $\rho_{ij} = \rho_{ik}\rho_{kj}$ and also $\rho_{ij} = \rho_{ji}^{-1}$. Over U_i we have $\mathcal{P}_{i,(F,\sigma)} \otimes \mathcal{P}_{i,(F,\sigma)} = \det(\widetilde{M}_i^\bullet)$. As usual, the $\det(\widetilde{M}_i^\bullet)$ path together (via $f_i\psi_i^{-1}$), to $\mathcal{D}_F$ and we get, again by Proposition 6.4, a canonical isomorphism $\mathcal{P}_{(F,\sigma)} \otimes \mathcal{P}_{(F,\sigma)} \xrightarrow{\sim} \mathcal{D}_F$. $\square$

(7.4) Considering the universal family over $\text{Bun}_{\text{SO}_r} \times X$, we get, by the above, for every theta-characteristic κ a line bundle $\mathcal{P}_\kappa$ over Bun_{SO_r} . Consider

$$e : \text{Bun}_{\text{Spin}_r} \longrightarrow \text{Bun}_{\text{SO}_r}(0)$$

defined by extension of the structure group. This morphism defines a morphism on the level of Picard groups hence we can define a line bundle, denoted by $\mathcal{P}$, which is the pullback of the pfaffian line bundle $\mathcal{P}_\kappa$. We omit here the index κ as we will see that $\mathcal{P}$ on $\text{Bun}_{\text{Spin}_r}$ does not depend on the choice of a particular theta-characteristic. In the same way, we define the line bundle $\mathcal{P}$ on Bun_{G_2} .

(7.5) *The pfaffian divisor.* Let $r \geq 3$ and $(\mathcal{E}, q)$ be the universal quadratic bundle over $\text{Bun}_{\text{SO}_r}(0) \times X$. For κ a theta-characteristic, let us denote by Θ_κ the substack defined by

$$\Theta_\kappa = \text{div}(Rpr_{1*}(\mathcal{E} \otimes pr_2^*\kappa)).$$

Claim: This substack is a divisor if and only if r or κ are even.

Proof. Let $P = (E, q)$ be a SO_r -bundle, $r \geq 3$ and κ be a theta-characteristic. Then

$$w_2(P) = h^0(E \otimes \kappa) + r h^0(\kappa) \bmod 2. \quad (7.5.1)$$

Indeed, by Riemann's invariance mod 2 theorem, the right hand side of (7.5.1) denoted $\bar{w}_2(P)$ in the following, is constant over the 2 connected components of Bun_{SO_r} . Because (7.5.1) is true at the trivial SO_r -bundle $\mathcal{T}$, it is enough to prove that $\bar{w}_2$ is not constant. Let $L, M \in J_2$ (where J_2 = points of order 2 of the jacobian) such that for the Weyl pairing $\langle L, M \rangle = 1$.

The choice of a trivialization of their square defines a non degenerated quadratic form on $E = (L \otimes M) \oplus L \oplus M \oplus (r-3)\mathcal{O}_X$ hence a SO_r -bundle P . By [23], we know that we have $\bar{w}_2(P) = \langle L, M \rangle \neq 0 = \bar{w}_2(\mathcal{T})$, which proves (7.5.1). Choose an ineffective theta-characteristic κ_0 and set $L = \kappa_0 \otimes \kappa^{-1}$. If r is even, there exists a SO_r -bundle $P = (E, q)$ such that $H^0(E \otimes \kappa) = 0$ and $w_2(P) = 0$ (choose $E = rL$ with $L \in J_2$ and use (7.5.1)). If r is odd and κ is even, there exists a SO_r -bundle $P = (E, q)$ such that $H^0(E \otimes \kappa) = 0$ and $w_2(P) = 0$ (by ([2], lemme 1.5), there is a SL_2 -bundle F on X such that $H^0(X, \mathrm{ad}(F) \otimes \kappa) = 0$, then choose $E = \mathrm{ad}(F) \oplus (r-3)L$ with the obvious quadratic form.) If r and κ are odd, then $H^0(E \otimes \kappa)$ is odd for all $P \in \mathrm{Bun}_{\mathrm{SO}_r}(0)$. $\square$

As the perfect complex $Rpr_{1*}(\mathcal{E} \otimes pr_2^* \kappa)$ can be locally represented by a skew-symmetric perfect complex of length one $L \xrightarrow{\alpha} L^*$, the pfaffian of α defines a local equation of an effective divisor $\frac{\Theta_\kappa}{2}$ such that $2\frac{\Theta_\kappa}{2} = \Theta_\kappa$. This gives an easier way to define, by smoothness of Bun_G , the pfaffian line bundle. The reason which motivated our construction above was to define this square root for arbitrary quadratic bundles (not only the even ones) and to make a construction for all theta-characteristics and not only the even ones (when r is odd).

(7.6) Proof of 1.5. The dualizing line bundle ω_{Bun_G} on Bun_G is by definition the determinant line bundle of the cotangent complex of Bun_G . Let $\mathrm{Ad} : G \rightarrow \mathrm{GL}(\mathfrak{g})$ be the adjoint representation. Then $\omega_{\mathrm{Bun}_G} = \mathcal{D}_{\mathrm{Ad}}^{-1}$. Suppose that G is semi-simple. Then the adjoint representation factors through the special orthogonal group because of the existence of the Cartan-Killing form. Choose a theta-characteristic κ on X . Then, as in (7.2), we can define a square root $\omega_{\mathrm{Bun}_G}^{\frac{1}{2}}(\kappa)$ of ω_{Bun_G} .

8. The Picard group of Bun_G .

Throughout the section suppose that $k = \mathbb{C}$ and that G is simple and simply connected.

Let $\mathrm{Pic}_{L_G^X}(\mathcal{Q}_G)$ the group of L_G^X -linearized line bundles on $\mathcal{Q}_G$. Recall that a L_G^X -linearization of the line bundle $\mathcal{L}$ on $\mathcal{Q}_G$ is an isomorphism $m^* \mathcal{L} \xrightarrow{\sim} pr_2^* \mathcal{L}$, where $m : L_G^X \times \mathcal{Q}_G \rightarrow \mathcal{Q}_G$ is the action of L_G^X on $\mathcal{Q}_G$, satisfying the usual cocycle condition.

Consider the projection $\pi : \mathcal{Q}_G \rightarrow \mathrm{Bun}_G$ of Theorem 1.3. Let $\mathcal{L}$ be a line bundle on Bun_G . As π^* induces an isomorphism between the sections of $\mathcal{L}$ and L_G^X -invariant sections of $\pi^* \mathcal{L}$ ([3], Lemma 7.2), we have

(8.1) PROPOSITION.— *The projection $\pi : \mathcal{Q}_G \rightarrow \mathrm{Bun}_G$ induces an injection*

$$\pi^* : \mathrm{Pic}(\mathrm{Bun}_G) \hookrightarrow \mathrm{Pic}_{L_G^X}(\mathcal{Q}_G).$$

Any L_G^X -linearization is necessarily unique:

(8.2) PROPOSITION.— *The forgetful morphism $\text{Pic}_{L_G^X}(\mathcal{Q}_G) \rightarrow \text{Pic}(\mathcal{Q}_G)$ is injective.*

Proof. The kernel of this morphism consists of the L_G^X -linearizations of the trivial bundle. Any two such trivializations differ by an automorphism of $pr_2^* \mathcal{O}_{\mathcal{Q}_G}$ that is by an invertible function on $L_G^X \times \mathcal{Q}_G$. Since $\mathcal{Q}_G$ is integral (4.2), it is the direct limit of the integral projective varieties $\mathcal{Q}_G = \varinjlim \mathcal{Q}_G(N)_{\text{red}}$ and this function is the pull back of an invertible function f on L_G^X . The cocycle conditions on the linearizations imply that f is a character, hence $f = 1$ by Lemma 5.2. $\square$

(8.3) $\text{Pic}(\mathcal{Q}(G))$ and the canonical central extension of L_G .

Consider the embedding $L_G/L_G^+ \hookrightarrow \mathbb{P}(\mathcal{H}_1)$ of (4.2) and define $\mathcal{O}_{\mathcal{Q}_G}(1)$ as the pullback of $\mathcal{O}_{\mathbb{P}(\mathcal{H}_1)}(1)$. By [21] and [17], we know $\text{Pic}(\mathcal{Q}_G) = \mathbb{Z}\mathcal{O}_{\mathcal{Q}_G}(1)$. The k -group L_G acts on $\mathcal{Q}_G$ but the action does not lift to an action of L_G on $\mathcal{O}_{\mathcal{Q}_G}(1)$. There is a canonical device to produce an extension $\widehat{L}_G$ of L_G such that the induced action of $\widehat{L}_G$ lifts to an action of $\widehat{L}_G$ on $\mathcal{O}_{\mathcal{Q}_G}(1)$: the Mumford group. This is the group of pairs (g, f) where $g \in L_G$ and $f : g^* \mathcal{O}_{\mathcal{Q}_G}(1) \xrightarrow{\sim} \mathcal{O}_{\mathcal{Q}_G}(1)$. We get a central extension (note that $\mathcal{Q}_G$ is direct limit of projective integral schemes)

$$1 \longrightarrow G_m \longrightarrow \widehat{L}_G \longrightarrow L_G \longrightarrow 1 \quad (8.3.1)$$

Note that the Mumford group of $\mathcal{O}_{\mathbb{P}(\mathcal{H}_1)}(1)$ is $\text{GL}(\mathcal{H}_1)$. As the projective representation $\mathcal{H}_1$ of $L_{\mathfrak{g}}$ can be integrated to a projective representation $\phi : L_G \rightarrow \text{PGL}(\mathcal{H}_1)$ (cf. 4.2), by functoriality of the Mumford group, (8.3.1) is also the pullback by ϕ of the central extension

$$1 \longrightarrow G_m \longrightarrow \text{GL}(\mathcal{H}_1) \longrightarrow \text{PGL}(\mathcal{H}_1) \longrightarrow 1. \quad (8.3.2)$$

Moreover, the restriction of $\mathcal{H}_1$ to $L_{\mathfrak{g}}^+$ can be integrated to a representation $L_G^+ \rightarrow \text{GL}(\mathcal{H}_1)$. It follows that (8.3.2) splits *canonically* over L_G^+ , and $\mathcal{O}_{\mathcal{Q}_G}(1)$ is the line bundle on the homogeneous space $\mathcal{Q}_G = \widehat{L}_G / L_G^+$ associated to the character $G_m \times L_G^+ \rightarrow G_m$ defined by the first projection.

(8.4) Consider the case of $G = \text{SL}_r$. In this case, we know from [3], that the pullback of the determinant line $\mathcal{D}$ bundle is $\mathcal{O}_{\mathcal{Q}_{\text{SL}_r}}(1)$. It follows from (8.1), (8.2) and (8.3) that $\text{Pic}(\text{Bun}_G) = \mathbb{Z}\mathcal{D}$. If $\rho : G \rightarrow \text{SL}_r$ is a representation of G we get a commutative diagram

$$\begin{array}{ccc} \mathcal{Q}_G & \longrightarrow & \mathcal{Q}_{\text{SL}_r} \\ \downarrow & \searrow \varphi & \downarrow \\ \text{Bun}_G & \longrightarrow & \text{Bun}_{\text{SL}_r} \end{array}$$

(8.5) LEMMA.— *Let d_ρ be the Dynkin index of ρ . Then the pullback of the determinant bundle under φ is $\mathcal{O}_{\mathcal{Q}_G}(d_\rho)$*

Proof. Consider the pullback diagram of (8.3.1) for LSL_r :

$$\begin{array}{ccccccccc} 1 & \longrightarrow & G_m & \longrightarrow & \widetilde{L}_G & \longrightarrow & L_G & \longrightarrow & 1 \\ & & \parallel & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & G_m & \longrightarrow & \widehat{\text{LSL}}_r & \longrightarrow & \text{LSL}_r & \longrightarrow & 1 \end{array}$$

Looking at the differentials (note that $\text{Lie}(\widehat{\text{LSL}}_r) = \widehat{L}_{\mathfrak{sl}_r}$ by [3]), on the level of Lie algebra, we restrict the universal central extension of $L_{\mathfrak{sl}_r}$ to $L_{\mathfrak{g}}$. The resulting extension $\widetilde{L}_{\mathfrak{g}}$ is (cf. Section (2.3)) the Lie algebra of the Mumford group of $\mathcal{O}_{\mathcal{Q}_G}(d_\rho)$ where d_ρ is the Dynkin index of d_ρ , which proves the lemma. $\square$

(8.6) COROLLARY.— *As a pullback, the line bundle $\mathcal{O}_{\mathcal{Q}_G}(d_\rho)$ is L_G^X -linearized.*

(8.7) Proof of Theorem 1.1. By the above for series A and C, as the Dynkin index for the standard representation is 1, that all line bundles on $\mathcal{Q}_G$ are L_G^X -linearized. For the series B and D (and also for G_2) the Dynkin index of the standard representation is 2. But by the existence of the pfaffian line bundles we see also in this case, that all line bundles on $\mathcal{Q}_G$ are L_G^X -linearized and Theorem 1.1 for $n = 0$ follows from (8.1), (8.2) and (8.3)

(8.8) REMARK.— The restriction to L_G^X of the canonical central extension $\widehat{L}_G$ of L_G splits (at least for classical G and G_2). Moreover this splitting is unique.

Proof. As $\mathcal{O}_{\mathcal{Q}_G}(1)$ admits a L_G^X -linearization, the action of L_G^X on $\mathcal{Q}_G$ induced by the embedding $L_G^X \subset L_G$ lifts to an action to $\mathcal{O}_{\mathcal{Q}_G}(1)$. The central extension of L_G^X obtained by pullback of the canonical central extension $\widehat{L}_G$ of L_G is the Mumford group of L_G^X associated to $\mathcal{O}_{\mathcal{Q}_G}(1)$. But this extension splits as the action lifts and we are done. Two splittings differ by a character of L_G^X . As there is only the trivial character (corollary 5.2) the splitting must be unique. $\square$

9. Parabolic G-bundles.

Throughout this section G is simple, simply connected and $k = \mathbb{C}$. We will extend the previous sections to the moduli stacks of parabolic G -bundles.

(9.1) We use the notations of Section 2. We will recall some standard facts for Lie groups, which we will use later. Let G be the simple and simply connected algebraic group associated to $\mathfrak{g}$. Denote by $T \subset G$ the Cartan subgroup associated to $\mathfrak{h} \subset \mathfrak{g}$ and by $B \subset G$ the Borel subgroup associated to $\mathfrak{b} \subset \mathfrak{g}$. Given a subset Σ of the set of simple roots Π (nodes of the Dynkin diagram), we can define a subalgebra $\mathfrak{p}_\Sigma = \mathfrak{b} \oplus (\oplus_{\alpha \in \Sigma} \mathfrak{g}_{-\alpha}) \subset \mathfrak{g}$, hence a subgroup $P_\Sigma \subset G$. Remark that $P_\emptyset = G/B$, $P_\Pi = G$ and that all P_Σ contain B . The subgroup P_Σ is parabolic and conversely any *standard* (i.e. containing B) parabolic subgroup arises in this way. Fix $\Sigma \subset \Pi$ and let $\Gamma = \Pi - \Sigma$. Denote by $X(P_\Sigma)$ the character group of P_Σ . Any weight λ such that $\lambda(H_\alpha) = 0$ for all $\alpha \in \Sigma$ defines, via the exponential map, a character of P_Σ and all characters arise in this way, i.e. $X(P_\Sigma) = \{\lambda \in P / \lambda(H_\alpha) = 0 \text{ for all } \alpha \in \Sigma\}$. Given $\chi \in X(P_\Sigma)$ we can define the line bundle $L_\chi = G \times^{P_\Sigma} k_\chi$ on the homogeneous space G/P_Σ . In general, there is an exact sequence ([11], prop. 3.1)

$$1 \longrightarrow X(G) \longrightarrow X(P_\Sigma) \longrightarrow \text{Pic}(G/P_\Sigma) \longrightarrow \text{Pic}(G) \longrightarrow \text{Pic}(P) \longrightarrow 0.$$

As G is simple, we have $X(G) = 0$ and as G is simply connected, we have $\text{Pic}(G) = 0$ ([11], cor. 4.5). We get the isomorphism $X(P_\Sigma) \xrightarrow{\sim} \text{Pic}(G/P_\Sigma)$. In particular, the Picard group of G/P_Σ is isomorphic to the free abelian group generated over Γ .

(9.2) Consider closed points $p_1, \dots, p_n$ of X , labeled with the standard parabolic subgroup $P_1, \dots, P_n$. Let $\Sigma_1, \dots, \Sigma_n$ be the associated subsets of simple roots and $\Gamma_i = \Pi - \Sigma_i$ for $i \in \{1, \dots, n\}$. In the following, underlining a character will mean that we consider the associated sequence, e.g. $\underline{P}$ will denote the sequence $(P_1, \dots, P_n)$, etc. Let E be a G -bundle. As G acts on G/P_i we can define the associated G/P_i -bundle $E(G/P_i)$.

(9.3) DEFINITION.— (cf. [22]) A *quasi-parabolic G -bundle of type $\underline{P}$* is a G -bundle E on X together with, for all $i \in \{1, \dots, n\}$, an element $F_i \in E(G/P_i)(p_i)$. A *parabolic G -bundle of type $(\underline{P}, \underline{m})$* is a quasi-parabolic G -bundle of type $\underline{P}$ together with, for $i \in \{1, \dots, n\}$, parabolic weights $(m_{i,j})_{j \in \Gamma_i}$ where the $m_{i,j}$ are strictly positive integers.

(9.4) Let R be a k -algebra, $S = \text{Spec}(R)$. A family of quasi-parabolic G -bundles of type $\underline{P}$ parameterized by S is a G -bundle E over $S \times X$ together with n sections $\sigma_i : S \rightarrow E(G/P_i)_{|S \times \{p_i\}}$. A morphism from $(E, \underline{\sigma})$ to $(E', \underline{\sigma}')$ is a morphism $f : E \rightarrow E'$ of G -bundles such that for all $i \in \{1, \dots, n\}$ we have $\sigma'_i = f_{|S \times \{p_i\}} \circ \sigma_i$.

We get a functor from the category of k -algebras to the category of groupoids by associating to R the groupoid having as objects families of quasi-parabolic G -bundles of type $\underline{P}$ parameterized by $S = \text{Spec}(R)$ and as arrows isomorphisms between such families. Moreover for any morphism $R \rightarrow R'$ we have a natural functor between the associated groupoids. This defines

the k -stack of quasi-parabolic G -bundles of type $\underline{P}$ which we will denote by $\text{Bun}_G^{par}(\underline{p}, \underline{P})$. The stack $\text{Bun}_G^{par}(\underline{p}, \underline{P})$ has, as Bun_G , a natural interpretation as a double quotient stack. Define

$$\mathcal{Q}_G^{par}(\underline{p}, \underline{P}) = \mathcal{Q}_G \times \prod_{i=1}^n G/P_i.$$

The ind-group L_G^X acts on $\mathcal{Q}_G$ and by evaluation $ev(p_i) : L_G^X \rightarrow G$ at p_i also on each factor G/P_i . We get a natural action of L_G^X on $\mathcal{Q}_G^{par}(\underline{p}, \underline{P})$. The analogue of Theorem 1.3. for quasi-parabolic G -bundles is

(9.5) THEOREM.— *(Uniformization) There is a canonical isomorphism of stacks*

$$\bar{\pi} : L_G^X \backslash \mathcal{Q}_G^{par}(\underline{p}, \underline{P}) \xrightarrow{\sim} \text{Bun}_G^{par}(\underline{p}, \underline{P}).$$

Moreover the projection map is locally trivial for the étale topology.

Proof. Let R be a k -algebra, $S = \text{Spec}(R)$. To an element $(E, \rho, \underline{f})$ of $\mathcal{Q}_G^{par}(\underline{p}, \underline{P})(R)$ (with $f_i \in \text{Mor}(S, G/P_i)$), we can associate a family of quasi-parabolic G -bundles of type $\underline{P}$ parameterized by S in the following way. We only have to define the sections σ_i :

$$\sigma_i : S \xrightarrow{(\text{id}, f_i)} S \times G/P_i \xrightarrow{\rho_i(G/P_i)} E(G/P_i)|_{S \times \{p_i\}}.$$

We get a L_G^X equivariant map $\pi : \mathcal{Q}_G^{par}(\underline{p}, \underline{P}) \rightarrow \text{Bun}_G^{par}(\underline{p}, \underline{P})$ which induces the map on the level of stacks $\bar{\pi} : L_G^X \backslash \mathcal{Q}_G^{par}(\underline{p}, \underline{P}) \rightarrow \text{Bun}_G^{par}(\underline{p}, \underline{P})$.

Conversely, let $(E, \underline{\sigma})$ be a family of quasi-parabolic G -bundles of type $\underline{P}$ parameterized by $S = \text{Spec}(R)$. For any R -algebra R' , let $T(R')$ be the set of trivializations ρ of $E_{R'}$ over $X_{R'}$. This defines a R -space T which by Theorem 3.12 is a L_G^X -torsor. To any element in $T(R')$, we can associate the family $\underline{f}$ by

$$f_i : S \xrightarrow{\sigma_i} E(G/P_i)|_{S \times \{p_i\}} \xrightarrow{\rho_i(G/P_i)^{-1}} S \times G/P_i \xrightarrow{pr_2} G/P_i.$$

In this way we associate functorially to objects $(E, \underline{\sigma})$ of $\text{Bun}_G^{par}(\underline{p}, \underline{P})(R)$ L_G^X -equivariant maps $\alpha : T \rightarrow \mathcal{Q}_G^{par}(\underline{p}, \underline{P})$. This defines a morphism of stacks

$$\text{Bun}_G^{par}(\underline{p}, \underline{P}) \longrightarrow L_G^X \backslash \mathcal{Q}_G^{par}(\underline{p}, \underline{P})$$

which is the inverse of $\bar{\pi}$. The second statement is clear from the proof of Theorem 1.3. $\square$

(9.6) We study first line bundles over $\mathcal{Q}_G^{par}(\underline{p}, \underline{P})$. Using (8.3), (9.1) and $H^1(G/P_i, \mathcal{O}) = 0$, we obtain the following proposition, proving, as L_G^X has no characters, Theorem 1.1.

(9.7) PROPOSITION.— *We have*

$$\text{Pic}(\mathcal{Q}_G^{par}(\underline{p}, \underline{P})) = \mathbb{Z}\mathcal{O}_{\mathcal{Q}_G}(1) \times \prod_{i=1}^n \text{Pic}(G/P_i) = \mathbb{Z}\mathcal{O}_{\mathcal{Q}_G}(1) \times \prod_{i=1}^n X(P_i).$$

10. Conformal blocs and generalized theta functions.

Throughout this section G is simple and simply connected and $k = \mathbb{C}$.

(10.1) Fix an integer $\ell \geq 0$ (the level) and let $p_1, \dots, p_n$ be distinct closed points of X (we allow $n = 0$ *i.e.* no points), each of it labeled with a dominant weight λ_i lying in the fundamental alcôve P_ℓ . Choose also another point $p \in X$, distinct from the points $p_1, \dots, p_n$. Define

$$\mathcal{H}_{\underline{\lambda}} = \mathcal{H}_\ell \otimes \left(\bigotimes_{i=1}^n L_{\lambda_i} \right).$$

and let $L_{\mathfrak{g}}^X$ be $\mathfrak{g} \otimes A_X$. We can map $L_{\mathfrak{g}}^X$ via the Laurent developpement at the point p to $L_{\mathfrak{g}}$. The restriction to $L_{\mathfrak{g}}^X$ of the universal central extension $\widehat{L_{\mathfrak{g}}}$ of $L_{\mathfrak{g}}$ splits by the residue theorem, hence $L_{\mathfrak{g}}^X$ may be considered as a *sub Lie-algebra* of $\widehat{L_{\mathfrak{g}}}$. In particular, $\mathcal{H}_\ell$ is a $L_{\mathfrak{g}}^X$ -module. Evaluating $X \otimes f \in L_{\mathfrak{g}}^X$ at the point p_i , we may consider L_{λ_i} as a $L_{\mathfrak{g}}^X$ -module. Therefore $\mathcal{H}_{\underline{\lambda}}$ is a (left) $L_{\mathfrak{g}}^X$ -module. Define the space of conformal blocks (or vacua) by

$$V_X(\underline{p}, \underline{\lambda}) = [\mathcal{H}_{\underline{\lambda}}^*]^{L_{\mathfrak{g}}^X} := \{ \psi \in \mathcal{H}_{\underline{\lambda}}^* / \psi(X \otimes f) = 0 \ \forall X \otimes f \in L_{\mathfrak{g}}^X \}.$$

This definition is Beauville's description [1] (see also [31]) of the space of conformal blocks of Tsuchiya, Ueno and Yamada [32].

The labeling of the points p_i induces $\Sigma_i = \{ \alpha \in \Pi / \lambda_i(H_\alpha) = 0 \}$, $\Gamma_i = \Pi - \Sigma_i$ and $m_{i,j} = \lambda_i(H_{\alpha_j})$ for $j \in \Gamma_i$, that is the type of a parabolic G -bundle. In particular we get, for $\ell \in \mathbb{N}$, a natural line bundle on the moduli stack $\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P})$ defined by

$$\mathcal{L}(\ell, \underline{m}) = \mathcal{L}^\ell \boxtimes \left(\bigotimes_{i=1}^n \left(\bigotimes_{j \in \Gamma_i} \mathbb{Z} \mathcal{L}_{i,j}^{m_{i,j}} \right) \right).$$

By construction, for the pull back of $\mathcal{L}(\ell, \underline{m})$ to $\mathcal{Q}_G^{\text{par}}(\underline{p}, \underline{P})$ we have

$$\pi^* \mathcal{L}(\ell, \underline{m}) = \mathcal{O}_{\mathcal{Q}_G}(\ell) \boxtimes \left(\bigotimes_{i=1}^n \mathcal{L}_{-\lambda_i} \right)$$

where $\mathcal{L}_{-\lambda_i}$ is the line bundle on the homogeneous space G/P_i defined by the character corresponding to the weight $-\lambda_i$.

(10.2) Proof of (1.2.1): We extend the method of [3] and [24].

Step 1: As a pullback, $\pi^* \mathcal{L}(\ell, \underline{m})$ is canonically L_G^X -linearized, that is equipped with $\varphi : m^*(\pi^* \mathcal{L}(\ell, \underline{m})) \xrightarrow{\sim} pr_2^*(\pi^* \mathcal{L}(\ell, \underline{m}))$. Denote by $[H^0(\mathcal{Q}_G^{\text{par}}(\underline{p}, \underline{P}), \pi^* \mathcal{L}(\ell, \underline{m}))]^{L_G^X}$ the space of L_G^X -invariant sections, that is the sections s such that $\varphi(m^*s) = pr_2^*s$. By Lemma 7.2 of [3] we have the canonical isomorphism

$$H^0(\text{Bun}_G^{\text{par}}(\underline{p}, \underline{P}), \mathcal{L}(\ell, \underline{m})) \xrightarrow{\sim} [H^0(\mathcal{Q}_G^{\text{par}}(\underline{p}, \underline{P}), \pi^* \mathcal{L}(\ell, \underline{m}))]^{L_G^X}$$

Denote by $[H^0(\mathcal{Q}_G^{par}(\underline{p}, \underline{P}), \pi^* \mathcal{L}(\ell, \underline{m}))]^{L_G^X}$ the sections annihilated by $\text{Lie}(L_G^X) = L_{\mathfrak{g}}^X$. By Proposition 7.4 of [3], using that L_G^X and $\mathcal{Q}_G^{par}(\underline{p}, \underline{P})$ are integral (5.1 and 4.2), we have the canonical isomorphism

$$[H^0(\mathcal{Q}_G^{par}(\underline{p}, \underline{P}), \pi^* \mathcal{L}(\ell, \underline{m}))]^{L_G^X} \xrightarrow{\sim} [H^0(\mathcal{Q}_G^{par}(\underline{p}, \underline{P}), \pi^* \mathcal{L}(\ell, \underline{m}))]^{L_{\mathfrak{g}}^X}$$

Step 2: By definition of $\widehat{L}_G$, the space $H^0(\mathcal{Q}_G^{par}(\underline{p}, \underline{P}), \pi^* \mathcal{L}(\ell, \underline{m}))$ is naturally a $\widehat{L}_G$ -module. Moreover we know that $\widehat{L}_G$ splits over L_G^X (at least for classical G and G_2) and that this splitting is *unique*. The action of $L_{\mathfrak{g}}^X \subset \widehat{L}_{\mathfrak{g}}$ deduced from this inclusion on $H^0(\mathcal{Q}_G^{par}(\underline{p}, \underline{P}), \pi^* \mathcal{L}(\ell, \underline{m}))$ is therefore the same as the preceding one.

Step 3: We have the canonical isomorphism of $\widehat{L}_G$ -modules

$$H^0(\mathcal{Q}_G^{par}(\underline{p}, \underline{P}), \pi^*(\mathcal{L}(\ell, \underline{m}))) \xrightarrow{\sim} H^0(\mathcal{Q}_G, \mathcal{O}_{\mathcal{Q}_G}(\ell)) \otimes \left(\bigotimes_{i=1}^n H^0(G/P_i, \mathcal{L}_{-\lambda_i}) \right)$$

To see this apply the Kunneth formula to the restriction of $\mathcal{L}(\ell, \underline{m})$ to the projective varieties $\mathcal{Q}_G^{par}(\underline{p}, \underline{P})^{(N)} = \mathcal{Q}_G(N) \times \prod_{i=1}^n G/P_i$, then use that inverse limits commute with the tensor products by finite dimensional vector spaces.

Step 4: We have the canonical isomorphism of $\widehat{L}_G$ -modules

$$H^0(\mathcal{Q}_G, \mathcal{O}_{\mathcal{Q}_G}(\ell)) \otimes \left(\bigotimes_{i=1}^n H^0(G/P_i, \mathcal{L}_{-\lambda_i}) \right) \xrightarrow{\sim} \mathcal{H}_{\ell,0}^* \otimes \left(\bigotimes_{i=1}^n L_{\lambda_i}^* \right)$$

This is Borel-Bott-Weil theory, in the version of Kumar-Mathieu ([16], [21]) for the first factor, and the standard version ¹ for the others.

The theorem follows from steps 1 to 4.

As we know the dimensions (at least for classical G and G_2) for the conformal blocks ([10],[1], or [31] for an overview) we get the Verlinde dimension formula for the spaces of generalized parabolic theta-functions.

11. Moduli spaces.

(11.1) Suppose $\text{char}(k) = 0$. We will show how the previous results apply to the *coarse moduli spaces* of principal G -bundles. We suppose that G is reductive and that $g \geq 2$. Recall that a G -bundle E over X is *semi-stable* (resp. *stable*) if for every parabolic subgroup P and for every reduction E_P of E to G , we have for every dominant character (with respect to some

¹In [5] only the case G/B (*i.e.* $\Sigma = \emptyset$) is considered but the generalization to arbitrary G/P_{Σ} is immediate (and well known)

Borel $B \subset P$) χ of P , trivial over $Z_0(G)$, the following inequality $\deg(E_P(\chi)) \leq 0$ (resp. $<$). A stable G -bundle E is called *regularly stable*, if moreover $\text{Aut}(E)/Z(G) = \{1\}$.

Topologically, G -bundles over X are classified by elements of $\pi_1(G)$. By Ramanathan's [26] theorem, there are coarse moduli spaces $\text{Bun}_G(\tau)$ of semi-stable principal G -bundles of dimension $(g-1)\dim G + \dim Z_0(G)$, which are irreducible, once the topological type $\tau \in \pi_1(G)$ is fixed. Moreover $M_G(\tau)$ is normal and the open subset $M_G^{\text{reg}}(\tau) \subset M_G(\tau)$ corresponding to regularly stable G -bundles is smooth.

(11.2) Denote Cl the group of Weil divisor classes. There is a commutative diagram

$$\begin{array}{ccc} \text{Pic}(M_G(\tau)) & \xrightarrow{c} & \text{Cl}(M_G(\tau)) \\ r_1 \downarrow & & \downarrow r_2 \\ \text{Pic}(M_G^{\text{reg}}(\tau)) & \xrightarrow{c_{\text{reg}}} & \text{Cl}(M_G^{\text{reg}}(\tau)) \end{array}$$

By normality, the restriction r_1 is injective, by smoothness of $M_G^{\text{reg}}(\tau)$, the canonical morphism c_{reg} is an isomorphism and as ([9], II.6) $\text{codim}_{M_G(\tau)} M_G(\tau) \setminus M_G^{\text{reg}}(\tau) \geq 2$ (except when $g = 2$ and G maps nontrivially to PGL_2) the restriction r_2 is an isomorphism. In particular, $M_G(\tau)$ is locally factorial [7] if and only if r_1 is surjective.

(11.3) Consider $G = \text{GL}_r$. Then we may present $M_G(0)$ as the good quotient $\mathcal{H}/\text{GL}(M)$ where $\mathcal{H}$ is Grothendieck's Quot scheme $\mathcal{H} = \text{Quot}^{ss}(k^M \otimes \mathcal{O}_X(-N), P)$ parameterizing equivalence classes (with the obvious equivalence relation) of pairs $[E, \alpha]$ with E a semi-stable vector-bundle of degree 0 and $\alpha : k^M \xrightarrow{\sim} E(N)$, where N and $M = rN + \chi(E)$. Let $\mathcal{E}$ be the universal family over $\mathcal{H} \times X$ and consider $D = \det(Rpr_{1*}(\mathcal{E} \otimes pr_2^*(L)))$, with L a line bundle. It is well known that $[E, \alpha] \in \mathcal{H}$ has closed orbit exactly when E is polystable, *i.e.* direct sum of stable bundles: $E \simeq E_1^{\oplus n_1} \oplus \dots \oplus E_\ell^{\oplus n_\ell}$, and that the action of the stabilizer $\text{GL}(n_1) \times \dots \times \text{GL}(n_\ell)$ is given by the character

$$(g_1, \dots, g_\ell) \mapsto \det(g_1)^{\chi(E_1 \otimes L)} \dots \det(g_\ell)^{\chi(E_\ell \otimes L)}.$$

Choose a line bundle L of degree $g-1$ on X . Then $\chi(E_q \otimes L) = 0$ for $q \in \mathcal{H}$ and the action is trivial. By Kempf's lemma [7], D descends to the determinant of cohomology line bundle on $M_{\text{GL}_r}(0)$.

(11.4) *Proof of 1.6.* Suppose G is simple and simply connected. We have (except for $g = 2$ and $G = \text{SL}_2$)

$$\text{codim}_{\text{Bun}_G}(\text{Bun}_G \setminus \text{Bun}_G^{\text{reg}}) \geq 2.$$

To see this define the Harder-Narasimhan filtration in the case of G -bundles and calculate the codimension of the strata ([20], Section 3) to show that for the open substack $\text{Bun}_G^{ss} \subset \text{Bun}_G$

corresponding to semi-stable G -bundles we have $\text{codim}_{\text{Bun}_G}(\text{Bun}_G \setminus \text{Bun}_G^{ss}) \geq 2$, then use ([9], II.6). The smoothness of Bun_G implies $\text{Pic}(\text{Bun}_G^{reg}) = \text{Pic}(\text{Bun}_G)$ and it follows from Theorem 1.1 that $\text{Pic}(M_G)$ is an infinite cyclic group (note that the canonical morphism $\text{Bun}_G^{reg} \rightarrow M_G^{reg}$ induces an injection on the level of Picard groups). By (11.3) and (2.3), we know that the generator is the determinant of cohomology for G of type A and C. Moreover, M_G is locally factorial by (11.2) in this case.

(11.5) Consider $G = \text{SO}_r$ with its standard (orthogonal) representation and suppose that $r \geq 7$. The moduli space $\text{Pic}(M_{\text{SO}_r})$ is the good quotient of a parameter scheme Quad by $\text{GL}(H)$ with $H = k^N$ (cf. [29]). The scheme Quad parameterizes equivalent (with the obvious equivalence relation) triples $([F, \sigma, \alpha])$, where (F, σ) is a semistable SO_r -bundle and $\alpha : H^0(X, F(N)) \xrightarrow{\sim} H$.

Choose a theta-characteristic κ on X . Then on Quad there is the $\text{GL}(H)$ -linearized pfaffian of cohomology line bundle $\mathcal{P}_\kappa$ deduced from the universal family over $\text{Quad} \times X$.

(11.6) PROPOSITION.— *The line bundle $\mathcal{P}_\kappa$ descends to $M_{\text{SO}_r}^{reg}$.*

Proof. We use Kempf's lemma. If r is even, the stabilizer at a point $q = [F, \sigma, \alpha] \in \text{Quad}^{reg}$ is ± 1 ; if r is odd, the stabilizer is reduced to 1. So in the latter case, there is nothing to prove. In the former case, by definition of the pfaffian of cohomology, using that its formation commutes with base change, the action ± 1 is given by $g \mapsto g^{h^1(F \otimes \kappa)}$, so the action is trivial, as $h^1(F \otimes \kappa)$ is even. $\square$

(11.7) PROPOSITION.— *If $r \geq 7$, the line bundle $\mathcal{P}_\kappa$ does not descend to $M_{\text{SO}_r}(0)$. In particular, $M_{\text{SO}_r}(0)$ is not locally factorial.*

Proof. Let (F_1, σ_1) be a regularly stable odd SO_4 -bundle, and (F_2, σ_2) be a regularly stable odd SO_{r-4} -bundle. If $r = 8$, suppose that (F_1, σ_1) and (F_2, σ_2) are not isomorphic. Then the orthogonal sum $(F, \tau) = (F_1 \oplus F_2, \sigma_1 \oplus \sigma_2)$ is even. Let $[F, \tau, \alpha] \in \text{Quad}$ be a point corresponding to (F, τ) . Again, by definition of the pfaffian of cohomology, using that its formation commutes with base change, we see that the action of the stabilizer $\{\pm 1\} \times \{\pm 1\}$ is

$$(g_1, g_2) \mapsto g_1^{h^1(F_1 \otimes \kappa)} g_2^{h^1(F_2 \otimes \kappa)}.$$

But then the element $(-1, 1)$ acts nontrivially. $\square$

References

- [1] A. BEAUVILLE *Conformal blocks, fusion rules and the Verlinde formula*, e-print alg-geom/9405001 (1994)
- [2] A. BEAUVILLE *Fibrés de rang deux sur une courbe, fibré déterminant et fonctions thêta, II*, Bull. Soc. math. France **119** (1991) 259-291
- [3] A. BEAUVILLE AND Y. LASZLO. *Conformal blocks and generalized theta functions*, Comm. Math. Physics **164** (1994) 385-419
- [4] A. BEAUVILLE AND Y. LASZLO. *A result on algebraic descent*, C.R.A.S (1995)
- [5] R. BOTT *Homogenous vector bundles*, Ann. of Math., Series 2 **66** (1957) 203-248
- [6] E. B. DYNKIN. *Semisimple subalgebras of semisimple Lie algebras*, AMS Tranl. Ser. II **6** (1957) 111-244
- [7] J. M. DREZET AND M. S. NARASIMHAN. *Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques*, Inv. math. **97** (1989) 53-94
- [8] V. DRINFELD AND C. SIMPSON. *B-structures on G-bundles and local triviality*, (To appear)
- [9] G. FALTINGS. *Stable G-bundles and projective connections*, J. Alg. Geometry **2** (1993) 507-568
- [10] G. FALTINGS. *A proof of the Verlinde formula*, J. Alg. Geometry **3** (1994) 347-374
- [11] FOSSUM AND IVERSEN. *On Picard groups of algebraic fiber spaces*, J. pure and appl. algebra **3** (1973) 269-280
- [12] G. HARDER. *Halbeinfache Gruppenschemata über Dedkindringen*, Inv. Math. **4** (1967) 165-191
- [13] V.G. KAC. *Infinite dimensional Lie algebras (Third edition)*, Cambride, Cambridge University Press (1990)
- [14] G. KEMPF. *Deformations of Semi-Euler Characteristics*, Amer. J. of Math., **114** (1992) 973-978

- [15] F. KNUDSEN AND D. MUMFORD. *The projectivity of the moduli space of stable curves I: preliminaries on “det” and “div”*, Math. Scand. **39** (1976) 19-55
- [16] S. KUMAR. *Demazure character formula in arbitrary Kac-Moody setting*, Inv. math. **89** (1987) 395-423
- [17] S. KUMAR, M. S. NARASIMHAN AND A. RAMANATHAN. *Infinite Grassmannians and Moduli Spaces of G-bundles*, Math. Annalen **300** (1994) 395-423
- [18] S. KUMAR AND M. S. NARASIMHAN. *Picard group of the moduli spaces of G-bundles* e-print alg-geom/9511012
- [19] G. LAUMON AND L. MORET-BAILLY. *Champs algébriques* Preprint Université Paris 11 (Orsay) (1992)
- [20] G. LAUMON AND M. RAPOPORT. *The Langlands lemma and the Betti numbers of stacks of G-bundles on a curve*, alg-geom/9503006 (1995)
- [21] O. MATHIEU. *Formules de caractères pour les algèbres de Kac-Moody générales*, Astérisque **159-160** (1988)
- [22] V.B. MEHTA AND C.S. SESHADRI. *Moduli of vector bundles on curves with parabolic structures*, Math. Annales **248** (1980) 205-239
- [23] D. MUMFORD. *Theta characteristics on an algebraic curve*, Ann. Sci. École Norm. Sup. **4** (1971) 181-192
- [24] C. PAULY. *Espaces de modules paraboliques et blocs conformes*, Preprint Université Paris 11 (Orsay) (1994)
- [25] S. RAMANAN AND A. RAMANTHAN. *Some remarks on the instability flag*, Tôhoku Math. Journal **36** (1984) 269-291
- [26] A. RAMANTHAN. *Stable Principal G-bundles*, PhD. Thesis (Bombay University) (1976)
- [27] I. R. ŠAFAREVIČ. *On some infinite-dimensional groups*, II Math. USSR-Izv. **18** (1982) 185-194
- [28] P. SLODOWY. *On the geometry of Schubert varieties attached to Kac-Moody Lie algebras*, Can. Math. Soc. Conf Proc. **6** (1984) 405-442

- [29] C. SORGER. *Thêta-caractéristiques des courbes tracées over une surface lisse*, J. reine angew. Math. **435** (1993) 83-118
- [30] C. SORGER. *La semi-caractéristique d'Euler-Poincaré des faisceaux ω -quadratiques sur un schéma de Cohen-Macaulay*, Bull. Soc. Math. France **122** (1994) 225-233
- [31] C. SORGER. *La formule de Verlinde*, Séminaire Bourbaki 794 (1994)
- [32] A. TSUCHIYA, K. UENO AND Y. YAMADA. *Conformal Field Theory on Universal Family of Stable Curves with Gauge Symmetries*, Adv. Studies in Pure Math. **19** (1989) 459-566

Yves LASZLO

DMI, ÉCOLE NORMALE SUPÉRIEURE, 45, RUE D'ULM, 75230 PARIS CEDEX 05, FRANCE
e-mail: yves.laszlo@ens.fr

Christoph SORGER

INSTITUT DE MATHÉMATIQUES JUSSIEU (UMR 9994 DU CNRS) - CASE POSTALE 7012,
 UNIVERSITÉ PARIS 7, 2, PLACE JUSSIEU, F-75251 PARIS CEDEX 05, FRANCE
e-mail: sorger@mathp7.jussieu.fr